

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 September 2002 (19.09.2002)

PCT

(10) International Publication Number
WO 02/072758 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number: PCT/US02/06912

(22) International Filing Date: 8 March 2002 (08.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/274,421 9 March 2001 (09.03.2001) US
60/275,597 13 March 2001 (13.03.2001) US

(71) Applicant: UNIVERSITY OF KENTUCKY RE-
SEARCH FOUNDATION [US/US]; A144, ASTECC
Building, Lexington, KY 40502 (US).

(72) Inventors: CHAPPELL, Joseph; 1808 Bimini Road,
Lexington, KY 40502 (US). RALSTON, Lyle, F.; 927
Quail Meadows Court, Chesterfield, MO 63017 (US).

(74) Agent: CLARK, Paul, T.; Clark & Elbing LLP, 101 Fed-
eral Street, Boston, MA 02110 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: CYTOCHROME P450S AND USES THEREOF

(57) Abstract: The invention features isolated cytochrome P450 polypeptides and nucleic acid molecules, as well as expression vectors and transgenic plants containing these molecules. In addition, the invention features uses of such molecules in methods of increasing the level of resistance against a disease caused by a plant pathogen in a transgenic plant, in methods for producing altered compounds, for example, hydroxylated compounds, and in methods of producing isoprenoid compounds.



WO 02/072758 A2

CYTOCHROME P450s AND USES THEREOF

5

Field of the Invention

This invention relates to cytochrome P450s and uses thereof.

Background of the Invention

10 Cytochrome P450s encompass a superfamily of oxidases responsible for the oxidation of numerous endobiotics and thousands of xenobiotics. In addition, in plants, cytochrome P450s play important roles in wound healing, pest resistance, signaling, and anti-microbial and anti-fungal activity.

 Capsidiol is a bicyclic, dihydroxylated sesquiterpene produced by many
15 Solanaceous species in response to a variety of environmental stimuli, including exposure to UV (Back et al., Plant Cell. Physiol. 389:899-904, 1998) and infection by microorganisms (Molot et al., Physiol. Plant Pathol. 379-389, 1981; Stolle et al., Phytopathology 78:1193-1197, 1988; Keller et al., Planta. 205:467-476, 1998). It is the primary antibiotic or phytoalexin produced in tobacco in response to fungal elicitation,
20 and it is derived from the isoprenoid pathway via its hydrocarbon precursor, 5-epi-aristolochene (Figure 1). Several of the biosynthetic enzymes leading up to 5-epi-aristolochene formation have been studied (Chappell, Annu. Rev. Plant Physiol. Plant Mol. Biol. 46:521-547, 1995), especially 5-epi-aristolochene synthase (EAS) (Vögeli and Chappell, Plant Physiol. 88:1291-1296, 1988; Back and Chappell, Proc. Natl. Acad. Sci. U.S.A. 93:6841-6845, 1996; Mathis et al., Biochemistry 36:8340-8348, 1997; Starks et al., Science 277:1815-1820, 1997). EAS commits carbon to sesquiterpene metabolism by catalyzing the cyclization of farnesyl diphosphate (FPP) to 5-epi-aristolochene. However, until the present invention, the enzyme(s) responsible for the conversion of 5-epi-aristolochene to capsidiol has yet to be fully identified and
25 characterized.
30

 Biochemical evidence from previous studies in tobacco (Whitehead et al., Phytochemistry 28:775-779, 1989) and green pepper (Hoshino et al., Phytochemistry 38:609-613, 1995) have suggested that the oxidation of 5-epi-aristolochene to capsidiol occurs in a two step process with one of the hydroxylation steps being constitutive and

the other being mediated by an elicitor-inducible cytochrome P450 (Figure 1). Because 1-deoxycapsidiol had been isolated from natural sources (Watson et al., Biochem. Soc. Trans. 11:589, 1983), Whitehead et al. (Phytochemistry 28:775-779, 1989), surmised that perhaps the biosynthesis of this intermediate was due to pathogen induction of a corresponding hydroxylase. They therefore prepared synthetic 1-deoxycapsidiol and reported a modest conversion of this compound to capsidiol when fed to control or unelicited tobacco cell cultures. This was further supported by their observation that radiolabeled 5-epi-aristolochene was only converted to capsidiol when fed to elicitor-induced cell cultures but not control cultures. Whitehead et al. (Phytochemistry 28:775-779, 1989) therefore concluded that the 3-hydroxylase, responsible for hydroxylation of 5-epi-aristolochene at C3 to generate 1-deoxycapsidiol, was pathogen/elicitor inducible, while the 1-hydroxylase, responsible for hydroxylating 1-deoxycapsidiol at the C1 to generate capsidiol, was constitutive. Hoshino et al. (Phytochemistry 38:609-613, 1995) added to the observations of Whitehead et al. (Phytochemistry 28:775-779, 1989) by directly measuring 3-hydroxylase-activity in microsomal preparations of arachidonic acid-elicited *Capsicum annuum* fruits and seedlings. These assays consisted of incubating 5-epi-aristolochene with microsome preparations and subsequently determining the amount of 1-deoxycapsidiol generated by a combination of thin-layer chromatography (TLC) separations and gas chromatography (GC). Their evidence demonstrated that the conversion of 5-epi-aristolochene to 1-deoxycapsidiol was dependent on both NADPH and O₂, and that 1-deoxycapsidiol accumulation *in vitro* was arrested by the P450 antagonists carbon monoxide (Omura and Sato, J. Biol. Chem. 239:2370-2378, 1964), ancymidol (Coolbaugh et al., Plant Physiol. 62:571-576, 1978), and ketoconazole (Rademacher, Annu. Rev. Plant Physiol. Plant Mol. Biol. 51:501-531, 2000).

Recent results suggest that the hydroxylation of 5-epi-aristolochene is an important regulated step in capsidiol biosynthesis. In studies to evaluate the effectiveness of methyl-jasmonate as an inducer of capsidiol biosynthesis in tobacco cell cultures, Mandujano-Chávez et al. (Arch. Biochem. Biophys. 381:285-294, 2000), reported that the modest accumulation of this phytoalexin was accompanied by a strong induction of EAS. This result implied that steps before or after the sesquiterpene cyclase reaction were limiting. Using an *in vivo* assay measuring the conversion rate of radiolabeled 5-epi-aristolochene to capsidiol, a very limited induction of the hydroxylase

activities was observed in cells treated with methyl jasmonate relative to that in fungal elicitor-treated cells. This result pointed to the hydroxylase reactions as a potentially limiting step in capsidiol biosynthesis.

5

Summary of the Invention

In one aspect, the invention features several isolated cytochrome P450 polypeptides (such as CYP71D20, CYP71D21, CYP73A27, CYP73A28, and CYP92A5, and P450s having substantial identity to these polypeptides), as well as isolated nucleic acid molecules that encode these P450s.

10

In related aspects, the invention features a vector (such as an expression vector) including an isolated nucleic acid molecule of the invention and a cell (for example, a prokaryotic cell, such as *Agrobacterium* or *E. coli*, or a eukaryotic cell, such as a mammalian, insect, yeast, or plant cell) including the isolated nucleic acid molecule or vector.

15

In yet another aspect, the invention features a transgenic plant or transgenic plant component including a nucleic acid molecule of the invention, wherein the nucleic acid molecule is expressed in the transgenic plant or the transgenic plant component.

Preferably, the transgenic plant or transgenic plant component is an angiosperm (for example, a monocot or dicot). In preferred embodiments, the transgenic plant or

20

transgenic plant component is a solanaceous, maize, rice, or cruciferous plant or a component thereof. The invention further includes a seed produced by the transgenic plant or transgenic plant component, or progeny thereof.

In another aspect, the invention features a method of providing an increased level of resistance against a disease caused by a plant pathogen in a transgenic plant. The

25

method involves: (a) producing a transgenic plant cell including the nucleic acid molecule of the invention integrated into the genome of the transgenic plant cell and positioned for expression in the plant cell; and (b) growing a transgenic plant from the plant cell wherein the nucleic acid molecule is expressed in the transgenic plant and the transgenic plant is thereby provided with an increased level of resistance against a

30

disease caused by a plant pathogen.

In another aspect, the invention features a method for producing an altered compound, the method including the steps of contacting the compound with one or more of the isolated polypeptides disclosed herein under conditions allowing for the

hydroxylation, oxidation, demethylation, or methylation of the compound and recovering the altered compound.

In still another aspect, the invention features a hydroxylating agent including any of the isolated polypeptides disclosed herein.

5 In yet another embodiment, the invention features an isolated nucleic acid molecule that specifically hybridizes under highly stringent conditions to the complement of any one of the sequences described in SEQ ID NO:2 (CYP71D20), SEQ ID NO:4 (CYP71D21), SEQ ID NO:6 (CYP73A27), SEQ ID NO:8 (CYP73A28), or SEQ ID NO:12 (CYP92A5), wherein such a nucleic acid molecule encodes a
10 cytochrome P450 polypeptide.

 In another aspect, the invention features a host cell expressing a recombinant isoprenoid synthase and a recombinant cytochrome P450. In preferred embodiments, the host cell further expresses, independently or in combination, a recombinant acetyltransferase, methyltransferase, or fatty acyltransferase. In other preferred
15 embodiments, the host expresses an endogenous or recombinant cytochrome reductase. Preferably, the host cell is a yeast cell, a bacterial cell, an insect cell, or a plant cell.

 In a related aspect, the invention features a method for producing an isoprenoid compound, the method including the steps of: (a) culturing a cell that expresses a recombinant isoprenoid synthase and a recombinant cytochrome P450 under conditions
20 wherein the isoprenoid synthase and the cytochrome P450 are expressed and catalyze the formation of an isoprenoid compound not normally produced by the cell; and (b) recovering the isoprenoid compound. In preferred embodiments, the host cell further expresses a recombinant acetyltransferase, a recombinant methyltransferase, or a recombinant fatty acyltransferase. In other preferred embodiments, the host cell
25 expresses an endogenous or recombinant cytochrome reductase. Preferably, the host cell is a yeast cell, a bacterial cell, an insect cell, or a plant cell.

 In yet another aspect, the invention features an isoprenoid compound produced according to the above-mentioned methods.

 By "P450 polypeptide," "cytochrome P450," or "P450" is meant a polypeptide
30 that contains a heme-binding domain and shows a CO absorption spectra peak at 450 nm according to standard methods, for example, those described herein. Such P450s may also include, without limitation, hydroxylase activity, dual hydroxylase activity,

demethylase activity, or oxidase activity. Such enzymatic activities are determined using methods well known in the art.

By "polypeptide" is meant any chain of amino acids, regardless of length or post-translational modification (for example, glycosylation or phosphorylation).

5 By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 80 or 85%, preferably 90%, more preferably 95%, and most preferably 97%, or even 98% identity to a reference amino acid sequence (for example, the amino acid sequence shown in SEQ ID NOS:1, 3, 5, 7, and 11) or nucleic acid sequence (for example, the nucleic acid sequences shown in SEQ ID NOS:2, 4, 6, 8, and 12,
10 respectively). For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110
15 nucleotides.

Sequence identity is typically measured using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705, BLAST, or PILEUP/PRETTYBOX programs). Such software matches identical
20 or similar sequences by assigning degrees of homology to various substitutions, deletions, and/or other modifications. Conservative substitutions typically include substitutions within the following groups: glycine alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

25 By an "isolated polypeptide" is meant a P450 polypeptide (for example, a CYP71D20 (SEQ ID NO: 1), CYP71D21 (SEQ ID NO:3), CYP73A27 (SEQ ID NO:5), CYP73A28 (SEQ ID NO:7), or CYP92A5 (SEQ ID NO:11) polypeptide) that has been separated from components that naturally accompany it. Typically, the polypeptide is isolated when it is at least 60%, by weight, free from the proteins and naturally-
30 occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, a P450 polypeptide. An isolated P450 polypeptide may be obtained, for example, by extraction from a natural source (for example, a plant cell); by

expression of a recombinant nucleic acid encoding a P450 polypeptide; or by chemically synthesizing the protein. Purity can be measured by any appropriate method, for example, column chromatography, polyacrylamide gel electrophoresis, or by HPLC analysis.

5 By "derived from" or "obtained from" is meant isolated from or having the sequence of a naturally-occurring sequence (e.g., cDNA, genomic DNA, synthetic, or combination thereof).

By "isolated nucleic acid molecule" is meant a nucleic acid molecule, e.g., a DNA molecule, that is free of the nucleic acid sequence(s) which, in the naturally-
10 occurring genome of the organism from which the nucleic acid molecule of the invention is derived, flank the nucleic acid molecule. The term therefore includes, for example, a recombinant DNA that is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule (for example, a cDNA or a genomic or cDNA
15 fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. The term "isolated nucleic acid molecule" also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence.

By "specifically hybridizes" is meant that a nucleic acid sequence is capable of hybridizing to a DNA sequence at least under low stringency conditions, and preferably
20 under high stringency conditions. For example, high stringency conditions may include hybridization at approximately 42°C in about 50% formamide, 0.1 mg/ml sheared salmon sperm DNA, 1% SDS, 2X SSC, 10% Dextran sulfate, a first wash at approximately 65°C in about 2X SSC, 1% SDS, followed by a second wash at approximately 65°C in about 0.1X SSC. Alternatively high stringency conditions may
25 include hybridization at approximately 42°C in about 50% formamide, 0.1 mg/ml sheared salmon sperm DNA, 0.5% SDS, 5X SSPE, 1X Denhardt's, followed by two washes at room temperature in 2X SSC, 0.1% SDS, and two washes at between 55-60°C in 0.2X SSC, 0.1% SDS. Reducing the stringency of the hybridization conditions may involve lowering the wash temperature and/or washing at a higher concentration of salt.
30 For example, low stringency conditions may include washing in 2X SSC, 0.1% SDS at 40°C.

By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding (as used herein) a P450 polypeptide.

By "positioned for expression" is meant that the DNA molecule is positioned
5 adjacent to a DNA sequence which directs transcription and translation of the sequence (i.e., facilitates the production of, for example, a P450 polypeptide, a recombinant protein, or an RNA molecule).

By "reporter gene" is meant a gene whose expression may be assayed; such genes include, without limitation, beta-glucuronidase (GUS), luciferase,
10 chloramphenicol transacetylase (CAT), green fluorescent protein (GFP), beta-galactosidase, herbicide resistant genes, and antibiotic resistance genes.

By "expression control region" is meant any minimal sequence sufficient to direct transcription. Included in the invention are promoter elements that are sufficient to render promoter-dependent gene expression controllable for cell-, tissue-, or organ-
15 specific gene expression, or elements that are inducible by external signals or agents (for example, light-, pathogen-, wound-, stress-, or hormone-inducible elements or chemical inducers such as salicylic acid (SA) or 2,2-dichloro isonicotinic acid (INA)); such elements may be located in the 5' or 3' regions of the native gene or engineered into a transgene construct.

By "operably linked" is meant that a gene and a regulatory sequence(s) are
20 connected in such a way as to permit gene expression when the appropriate molecules (for example, transcriptional activator proteins) are bound to the regulatory sequence(s).

By "plant cell" is meant any self-propagating cell bounded by a semi-permeable membrane and typically is one containing a plastid. Such a cell also requires a cell wall
25 if further propagation is desired. Plant cell, as used herein includes, without limitation, algae, cyanobacteria, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores.

By "plant component" is meant a part, segment, or organ obtained from an intact plant or plant cell. Exemplary plant components include, without limitation, somatic
30 embryos, leaves, stems, roots, flowers, tendrils, fruits, scions, and rootstocks.

By "transgene" is meant any piece of DNA which is inserted by artifice into a cell and typically becomes part of the genome, for example, the nuclear or plastidic genome, of the organism which develops from that cell. Such a transgene may include a

gene which is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

By "transgenic" is meant any cell which includes a DNA sequence which is inserted by artifice into a cell and becomes part of the genome of the organism which
5 develops from that cell. As used herein, the transgenic organisms are generally transgenic plants and the DNA (transgene) is inserted by artifice into the nuclear or plastidic genome. A transgenic plant according to the invention may contain one or more engineered traits.

By "pathogen" is meant an organism whose infection of viable plant tissue elicits
10 a disease response in the plant tissue. Such pathogens include, without limitation, bacteria, mycoplasmas, fungi, insects, nematodes, viruses, and viroids. Plant diseases caused by these pathogens are described in Chapters 11-16 of Agrios, Plant Pathology, 3rd ed., Academic Press, Inc., New York, 1988.

By "increased level of resistance" is meant a greater level of resistance to a
15 disease-causing pathogen in a transgenic plant (or cell or seed thereof) of the invention than the level of resistance relative to a control plant (for example, a non-transgenic plant). In preferred embodiments, the level of resistance in a transgenic plant of the invention is at least 20% (and preferably 30% or 40%) greater than the resistance of a control plant. In other preferred embodiments, the level of resistance to a disease-
20 causing pathogen is 50% greater, 60% greater, and more preferably even 75% or 90% greater than a control plant; with up to 100% above the level of resistance as compared to a control plant being most preferred. The level of resistance is measured using conventional methods. For example, the level of resistance to a pathogen may be determined by comparing physical features and characteristics (for example, plant height
25 and weight, or by comparing disease symptoms, for example, delayed lesion development, reduced lesion size, leaf wilting and curling, water-soaked spots, and discoloration of cells) of transgenic plants.

By "purified antibody" is meant antibody which is at least 60%, by weight, free from proteins and naturally-occurring organic molecules with which it is naturally
30 associated. Preferably, the preparation is at least 75%, more preferably 90%, and most preferably at least 99%, by weight, antibody, for example, an acquired resistance polypeptide-specific antibody. A purified P450 antibody may be obtained, for example,

by affinity chromatography using a recombinantly-produced P450 polypeptide and standard techniques.

By “specifically binds” is meant an antibody which recognizes and binds a P450 protein but which does not substantially recognize and bind other molecules in a sample, for example, a biological sample, which naturally includes a P450 protein such as CYP71D20, CYP71D21, CYP73A27, CYP73A28, or CYP92A5.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Brief Description of the Drawings

Figure 1 is a schematic diagram of a proposed alternative pathway for the biosynthesis of capsidiol in elicitor-treated *Nicotiana tabacum* cells. 5-epi-aristolochene is synthesized from FPP by the action of a sesquiterpene cyclase, 5-epi-aristolochene synthase (EAS), and is subsequently hydroxylated at C1 and C3 to form capsidiol.

Figure 2 is a graph showing an induction time course for sesquiterpene cyclase enzyme activity and sesquiterpene hydroxylase activity in cellulase-treated cell cultures. Sesquiterpene cyclase (5-epi-aristolochene synthase, EAS) enzyme activity was determined in extracts prepared from control (open squares) and elicitor-treated (closed squares) cells collected at the indicated time points. Sesquiterpene hydroxylase activity was determined using an indirect assay for control (open circles) and elicitor-treated (closed circles) cells. Cell cultures were incubated with [³H]-5-epi-aristolochene for 3 hours ending at the indicated time points before quantifying the incorporation of radioactivity into extracellular capsidiol, a dihydroxylated form of aristolochene (Mandujano-Chávez et al., Arch. Biochem. Biophys. 381:285-294, 2000).

Figure 3 is a series of graphs showing the dose dependent inhibition of 5-epi-aristolochene hydroxylase activity by ancymidol and ketoconazole. Cell cultures were incubated in the presence of cellulase (0.5 µg/mL) plus the indicated concentrations of ancymidol (A) or ketoconazole (B) for 12 hours prior to measuring the *in vivo* 5-epi-aristolochene hydroxylase activity in the cell suspension cultures (squares), or the EAS enzyme activity in extracts prepared from the collected cells (triangles). The *in vitro* activity of a purified EAS preparation (Back and Chappell, J. Biol. Chem. 270:7375-7381, 1995) was also measured at the indicated inhibitor concentrations as an additional test for non-specific effects of these inhibitors (circles).

Figure 4A is a schematic diagram of the primary structure of a generalized cytochrome P450 with conserved domains used for the design of PCR primers highlighted (SEQ ID NOS:26-29).

Figure 4B is a list of the degenerate P450-specific primers (SEQ ID NOS:30-36) that were used in various combinations with vector specific primers in the amplification of cytochrome P450 cDNA fragments.

Figure 4C is a scanned image of an ethidium bromide-stained agarose gel showing the PCR products amplified from a directional cDNA library prepared with mRNA isolated from elicitor-treated cells using the degenerate primer GRRXCP(A/G)-for (SEQ ID NO:35) and the T7 vector-specific primer (SEQ ID NO:37). The T3 vector-specific primer is also shown (SEQ ID NO:38).

Figure 5 is a series of Northern blots showing the induction time course for CYP71D, CYP73A, CYP82E, CYP92A, and EAS transcript accumulation in elicitor treated cells. Total RNA was extracted from tobacco suspension cells incubated with the cellulase elicitor for the indicated durations, size fractionated by agarose gel electrophoresis under denaturing conditions, and transferred to a nylon membrane before probing with the respective full-length cDNAs. The uniformity of sample loading was verified by ethidium bromide staining of ribosomal RNA (Loading control).

Figure 6 is a series of graphs showing carbon monoxide (CO) difference spectra of the microsomal fraction isolated from yeast expressing the CYP92A5 (A) and CYP71D20 (B) cDNAs. Expression of the respective plasmid constructs engineered into the yeast (WAT11) cells was induced by a galactose treatment, followed by isolation of microsomal preparations. The difference adsorption spectra of microsomes incubated in the presence (solid lines) and absence (broken lines) of carbon monoxide was determined.

Figure 7 is a series of gas chromatograms of the reaction products formed upon incubation of microsomes isolated from WAT11 yeast cells containing the CYP71D20 expression construct (A and C) or vector control DNA (B and D) with sesquiterpene substrates. Microsomes isolated from the indicated yeast lines were incubated with 5-epi-aristolochene (A and B) or 1-deoxycapsidiol (C and D) in the presence (solid lines) or absence (dashed lines) of NADPH. The identities of 5-epi-aristolochene, 1-deoxycapsidiol, and capsidiol were verified by mass spectrometry.

Figure 8 is a sequence comparison of the amino acid sequence of *Nicotiana tabacum* 5-epi-aristolochene (sesquiterpene) hydroxylase NtCYP71D20 (SEQ ID NO:1) with other plant terpene hydroxylases (SEQ ID NOS:39-43). NrCYP71A5v1 (GenBank accession number CAA70575) catalyzes the mono-hydroxylation of nerol and geraniol, linear monoterpenes, while PaCYP71A1 (A35867) catalyzes the epoxidation of these substrates (Hallahan et al., Biochim. Biophys. Acta. 1201:94-100, 1994). MsCYP71D18 (AAD44150) and MpCYP71D13 (AAD44151) catalyze the mono-hydroxylation at C6 and C3 of limonene, a cyclic monoterpene, respectively (Lupien et al., Arch. Biochem. Biophys. 368:181-192, 1999). AtCYP701A3 (AAC39505) encodes for kaurene oxidase, which catalyzes a 3-step reaction including a hydroxylation followed by oxidation of a diterpene (Helliwell et al., Plant Physiol. 119:507-510, 1999). Shown are sequences from *Mentha piperita* (MpCYP71D13; SEQ ID NO:39), *Mentha spicata* (MsCYP71D18; SEQ ID NO:40), *Nepeta racemosa* (NrCYP71A5v1; SEQ ID NO:41), *Nicotiana tabacum* (NtCYP71D20; SEQ ID NO:1), *Persea americana* (PaCYP71A1; SEQ ID NO:42), and *Arabidopsis thaliana* (CYP701A3; SEQ ID NO:43). Conserved residues are shaded.

Detailed Description

Capsidiol is a bicyclic, dihydroxylated sesquiterpene produced by several Solanaceous species in response to a variety of environmental stimuli. It is the primary antimicrobial compound produced by *Nicotiana tabacum* in response to fungal elicitation, and it is formed via the isoprenoid pathway from 5-epi-aristolochene. Much of the biosynthetic pathway for the formation of this compound has been elucidated, except for the enzyme(s) responsible for the conversion of the allylic sesquiterpene 5-epi-aristolochene to its dihydroxylated form, capsidiol.

Accordingly, an *in vivo* assay for 5-epi-aristolochene hydroxylase-activity was developed and used to demonstrate a dose dependent inhibition of activity by ancyimidol and ketoconazole, two well-characterized inhibitors of cytochrome P450 enzymes. Using degenerate oligonucleotide primers designed to the well-conserved domains found within most P450 enzymes, including the heme binding domain, cDNA fragments representing four distinct P450 families (CYP71, CYP73, CYP82, and CYP92) were amplified from a cDNA library prepared against mRNA from elicitor-treated cells using PCR. The PCR fragments were subsequently used to isolate full-length cDNAs

(CYP71D20 (SEQ ID NO: 2) and D21 (SEQ ID NO: 4), CYP73A27 (SEQ ID NO: 6) and A28 (SEQ ID NO: 8), CYP82E1 (SEQ ID NO: 10), and CYP92A5 (SEQ ID NO: 12)), and these in turn were used to demonstrate that the corresponding mRNAs were all induced in elicitor-treated cells, albeit with different induction patterns.

5 There now follows a description of the cloning of several P450s from *Nicotiana tabacum*. These examples are provided for the purpose of illustrating the invention, and are not to be considered as limiting.

Inhibition of the 5-epi-aristolochene to capsidiol conversion by P450 antagonists

10 Using an indirect assay, a detailed induction time course of 5EAH activity in elicitor-induced cell cultures was determined relative to that of EAS activity (Figure 2), the well-characterized sesquiterpene cyclase activity that catalyzes the formation of 5-epi-aristolochene from FPP (Figure 1). Using assays for EAS and 5EAH, EAS activity is not detectable in control cell cultures, but is induced significantly within 3 hours and
15 reaches its maximal level within 15 to 18 hours of elicitor-treatment. Similar to the EAS enzyme activity, 5EAH activity was negligible in control cell cultures. Nonetheless, after an apparent lag phase of 8 hours, a rapid induction of hydroxylase activity was observed 10 to 15 hours post elicitor addition to the cell cultures, reaching a maximum by 18 hours followed by a rather gradual decline of 10 to 20% over the next 8 hours.

20 Tobacco cell suspension cultures treated with cellulase plus varying concentrations of ancymidol or ketoconazole were pre-incubated for 12 hours before measuring the cells' ability to convert exogenous supplied [³H] labeled 5-epi-aristolochene to radiolabeled capsidiol during a subsequent 3 hour incubation period (Figure 3). Apparent activity of 5EAH was inhibited in a dose-dependent manner with
25 approximately 50% inhibition by either 25 μ M ancymidol or ketoconazole, and more than 80% by 75 μ M ancymidol and 95% by 100 μ M ketoconazole (Figure 3A and B). Importantly, neither the *in vitro* activity of recombinant EAS nor the induction of EAS in the elicitor-treated cell cultures was significantly affected by ancymidol at concentrations as high as 100 μ M (Figure 3A). Ketoconazole also does not appear to
30 affect the *in vitro* activity of EAS. However, the inducibility of cyclase activity in elicitor-treated cell extracts was inhibited by ketoconazole at concentrations above 50 μ M (Figure 3B). Therefore, the specificity of ketoconazole as an inhibitor of P450 type

reactions should be assessed at or below a concentration of 50 μ M under these experimental conditions.

Isolation of elicitor-inducible cytochrome P450 cDNAs

5 A two-step approach for the isolation of candidate P450 cDNAs was followed. A PCR strategy was first employed using a directional cDNA library prepared against mRNA isolated from elicitor-induced cells as the template and degenerate PCR primers (Figure 4). Sequence alignments of cytochrome P450s from multiple families across
10 kingdoms were used to identify conserved regions to which a series of degenerate primers were prepared (Figure 4A and B). In cloning experiments, 450 to 550 bp products were expected from reactions utilizing the primer prepared to the heme-binding domain (GRRXCP(A/G)) (SEQ ID NOS:27 and 28) and the T7 vector primer (Figure 4C). The mixtures of reaction products were shotgun cloned, and approximately 100 of the cloned PCR fragments were sequenced. About half of the sequenced DNAs
15 contained signature sequences typical of P450 enzymes as revealed by BlastX database searches, and these corresponded to typical plant P450 family members of the CYP71, CYP73, CYP92, and CYP82 classes. Each of these PCR fragments was isolated multiple times in separate experiments. In addition, we isolated full-length cDNAs for these P450 family members. Table 1 compares the similarity and identity of the full-
20 length cDNAs of P450 family members with those of their nearest family member in the GenBank database. In addition, Figure 8 shows an amino acid alignment of several terpene cytochrome P450s. Alignments were performed using the algorithm of the MACVECTOR software suite.

25

Table 1: Full-length cDNAs cloned from an elicited cDNA library

Cytochrome P450 cDNA clone	Nearest relative/ accession number	% identity	% similarity
CYP71D20	CYP71D7 (<i>S. chacoense</i>) Gen EMBL U48435	76.5	88.8
CYP71D21	CYP71D7 (<i>S. chacoense</i>) Gen EMBL U48435	76.3	88.8
CYP73A27	CYP73A15 (<i>P. vulgaris</i>) Gen EMBL Y09447	79.4	92.6
CYP73A28	CYP73A15 (<i>P. vulgaris</i>) Gen EMBL Y09447	79.2	92.4
CYP82E1	CYP82E1 (<i>N. tabacum</i>) Gen EMBL AB015762	100.0	100.0
CYP92A5	CYP92A3 (<i>N. tabacum</i>) Gen EMBL X96784	95.5	98.6

The cloned fragments were used in a second step to isolate full-length clones from the cDNA library. Screening the cDNA library by hybridization with the CYP71 and CYP73 gene fragments yielded four full-length cDNAs, two CYP71Ds and two CYP73As. The former clones were designated CYP71D20 and CYP71D21, and the latter were designated CYP73A27 and CYP73A28. The other two cDNA fragments corresponded to tobacco cDNAs already found in the GenBank database, CYP82E1 and CYP92A3. These two cDNAs were cloned using specific primers designed with the help of the available sequence information to amplify the full-length cDNA

Induction of cytochrome P450 mRNAs in elicitor-treated cells

To correlate a biochemical role for P450s in sesquiterpene metabolism, RNA blot analyses were used to determine the steady-state levels of the mRNAs coding for all four of the cytochrome P450 clones and EAS in control and elicitor-treated cells (Figure 5). The mRNAs for all four of the P450s were rapidly and transiently induced with slightly different time courses relative to one another and to the EAS mRNA. CYP73A27 mRNA, for instance, displayed an induction pattern similar to that of EAS

with the maximum mRNA level occurring 9 to 12 hours after elicitation. While the EAS mRNA remained high throughout the duration of the experiment, the CYP73A27 mRNA was negligible in cells 24 hours after elicitor-treatment. In contrast, the CYP71D mRNA was more rapidly induced than the EAS mRNA, reached its maximum 6 to 9 hours after elicitation, and was declining by 12 hours when the EAS mRNA level was still very high.

Functional identification of CYP71D20 as 5-epi-aristolochene hydroxylase

To ascribe functional identity to the various P450 cDNAs, full-length cDNAs for CYP71D20, CYP82E1 and CYP92A5 were inserted into the yeast expression vector pYeDP60 (Urban et al., *Biochimie* 72:463-472, 1990; Pompon et al., *Methods Enzymol.* 272:51-64, 1996) and the expression of each in WAT11, a yeast line containing an integrated *Arabidopsis thaliana* cytochrome reductase gene (Pompon et al., *Methods Enzymol.* 272:51-64, 1996; Urban et al., *J. Biol. Chem.* 272:19176-19186, 1997), was determined. Engineering the CYP73A27 cDNA required an extra modification because of an unusually long N-terminus with several hydrophilic residues that may interfere with proper intracellular targeting (Nedelkina et al., *Plant Mol. Biol.* 39:1079-1090, 1999). This unusual leader sequence therefore was replaced with the membrane anchoring sequence of CYP73A1, a cinnamate 4-hydroxylase previously demonstrated to express well in yeast (Fahrendorf and Dixon, *Arch. Biochem. Biophys.* 305:509-515, 1993; Pompon et al., *Methods Enzymol.* 272:51-64, 1996). Expression of all these cDNAs was under the control of the glucose-repressible, galactose-inducible GAL10-CYC1 promoter (Guarente et al., *Proc. Natl. Acad. Sci. U.S.A.* 79:7410-7414, 1982), and expression was compared to yeast transformed with the parent pYeDP60 vector (control) alone.

After induction with galactose for approximately 16 hours, control cells and cells containing the various P450 constructs were collected, and microsomes prepared from each were analyzed for general P450 expression by CO-difference spectroscopy (Omura and Sato, *J. Biol. Chem.* 239:2370-2378, 1964). Microsomes prepared from cells containing the CYP71D20 (Figure 6A) and CYP92A5 (Figure 6B) constructs both showed characteristic CO difference spectra with peaks at 450 nm, indicating that the encoded proteins were assembling properly with their heme cofactor. Using the extinction coefficient of $91 \text{ mM}^{-1} \cdot \text{cm}^{-1}$ for heme binding proteins (Omura and Sato, J.

Biol. Chem. 239:2370-2378, 1964), it was determined that approximately 107 pmol of CYP71D20 and 268 pmol of CYP92A5 were expressed in the yeast cells per milligram of total yeast protein.

Both 5-epi-aristolochene and 1-deoxycapsidiol were metabolized to only one product with the same retention time as capsidiol. Obvious by its absence, no reaction product having a retention time similar to deoxycapsidiol was detectable in the 5-epi-aristolochene incubations (Figure 7). Co-injection of authentic capsidiol with the respective reaction products resulted in a single GC peak having a 16.2 minute retention time, identical to capsidiol. Mass spectra patterns for the separate reaction products were identical to that for the capsidiol standard (EIMS m/z 236, 221, 203, 185, 175, 163, 157, 133, 121, 107, 93, 79, 67, 55, 43, 41).

The *in vivo* assay data presented in Figures 2 and 3 of the current work indicate that the conversion of 5-epi-aristolochene is catalyzed by at least one inducible cytochrome P450 mediated reaction.

Furthermore, any of the cytochrome p450 polypeptides described herein may include one or more hydroxylase activities which can incorporate hydroxyl groups into at least two distant sites on an isoprenoid compound. The addition of these hydroxyl groups may occur, for example, sequentially, by adding a hydroxyl group first to one site and then the other, in either order. Moreover, such hydroxylases may be mutated to limit their ability to hydroxylate a substrate at only one site, or, alternatively, to provide stereochemical specificity to their hydroxylating activity.

The above-described experiments were performed using the following materials and methods.

Chemicals

Standard laboratory reagents were purchased from Becton Dickinson Microbiology Systems (Sparks, MD), FisherBiotech (Fair Lawn, NJ) and Sigma Chemical Company (St. Louis, MO).

Biological materials and induction treatments

Nicotiana tabacum cv. KY14 plants and cell suspension cultures were used. Cell suspension cultures were maintained in modified Murashige-Skoog (Vögeli and Chappell, Plant Physiol. 88:1291-1296, 1988). Cultures in their rapid phase of growth

(3 days old) were used for all experiments. At the indicated times, cells were collected and separated from media by vacuum filtration and stored at -80°C.

Induction treatments were performed by the addition of the fungal elicitors, cellulase (*Trichoderma viride*, Type RS, Onozuka) or paraciticein (O'Donohue et al.,
5 Plant Mol. Biol. 27:577-586, 1995) at the indicated concentrations. Paraciticein was purified from *E. coli* cells overexpressing a recombinant paraciticein protein containing a carboxy-terminal histidine purification tag.

In vivo 5-epi-aristolochene hydroxylase assay and inhibition studies

10 5-epi-aristolochene hydroxylase-activity was measured as the incorporation of [³H]-5-epi-aristolochene into extracellular capsidiol by intact cells. [³H]-5-epi-aristolochene was produced by incubating an excess of [1-³H] farnesyl diphosphate (1 μM, 20.5 Ci/mmol) with recombinant 5-epi-aristolochene synthase (Back et al., Arch. Biochem. Biophys. 315:527-532, 1994; Rising et al., J. Am. Chem. Soc. 122:1861-1866,
15 2000). The hexane extractable radioactivity from reactions was treated with a small amount of silica to remove any farnesol or residual FPP before quantifying the yield of radioactive 5-epi-aristolochene by liquid scintillation counting. The hexane solvent was removed under a gentle stream of N₂ gas, and the dried residue was re-dissolved in acetone. Control and elicitor-treated cells were then incubated with [³H]-5-epi-
20 aristolochene (approximately 100,000 dpm at 2.5 nM) for 3 hour periods at various points during an induction time course before collecting the cell and media samples. Detection and quantification of capsidiol in the extracellular culture media was performed as reported previously (Chappell et al., Phytochemistry 26:2259-2260, 1987), and the amount of radioactivity incorporated into capsidiol was determined. For these
25 determinations, samples were separated by TLC, and the zones corresponding to capsidiol were scraped from the plate for scintillation counting.

Inhibition studies were performed by the addition of the P450 inhibitors ancymidol (Coolbaugh et al., Plant Physiol. 62:571-576, 1978; Hoshino et al., Phytochemistry 38:609-613, 1995) and ketoconazole (Hoshino et al., Phytochemistry
30 38:609-613, 1995; Rademacher, Annu. Rev. Plant Physiol. Plant Mol. Biol. 51:501-531, 2000) directly to the cell cultures or enzyme assay mix. Cell cultures were incubated in the presence of cellulase (0.5 μg/ml) and indicated concentrations of ancymidol or ketoconazole for 12 hours prior to the addition of [³H]-5-epi-aristolochene. After a

further 3 hour incubation period, the cells and media were collected. The amount of radioactivity incorporated into extracellular capsidiol was determined as described above. To evaluate secondary effects of these inhibitors, the level of inducible sesquiterpene cyclase activity in the collected cells was determined according to (Vögeli et al., Plant Physiol. 93:182-187, 1990), as well as *in vitro* assays with purified recombinant EAS (Back et al., Arch. Biochem. Biophys. 315:527-532, 1994) incubated with the indicated concentrations of ancyimidol and ketoconazole.

All experiments were replicated in several independent trials. While the absolute values presented may have varied between experiments by as much as 50%, the trends and time courses were consistent throughout.

Construction of an elicitor-induced cDNA library

Cell cultures were incubated with fungal elicitor (0.5 µg cellulase/ml) for 6 hours before collecting the cells by filtration. The cells were kept frozen at -80°C until total RNA was extracted from them using Trizol (Life Technologies, Rockville, MD) according to the manufacturer's instructions. Poly.(A)⁺ RNA was purified by two rounds of oligo (dT) cellulose column chromatography (Life Technologies, Rockville, MD). cDNA synthesis and library construction were subsequently carried out using the UNI-ZAP XR library kit (Stratagene, La Jolla, CA), according to manufacturer's instructions.

PCR cloning strategy

Cytochrome P450 cDNA fragments were amplified from the elicitor-induced cDNA library using various combinations of degenerate forward and reverse primers with the vector-specific T3 and T7 primers. The template DNA was prepared from a 500 µl aliquot of the elicitor-induced cDNA library (3×10^6 pfu/µl) by heat denaturation at 70°C for 10 minutes, followed by phenol/chloroform extraction, ethanol precipitation and re-suspension in 500 µl of sterile, deionized water. Amplification reactions were performed in 50 µl volumes containing 50 mM KCl; 10 mM Tris-HCl, pH 8.8; 1.5 mM MgCl₂; 200 µM of each dNTP; 2 µl template DNA; 20 pmol each of forward and reverse primer; and 1 unit Taq Polymerase (Life Technologies, Rockville, MD). Reactions were preheated at 94°C for 2 minutes, followed by thirty-five cycles of denaturing at 94°C for 1 minute, annealing at 50°C for 1 minute 30 seconds, and

polymerization at 72°C for 2 minutes. The reactions were completed by a 10-minute extension at 72°C. Aliquots of the reaction products were examined for DNA products by agarose gel fractionation, and ligated directly into the pGEM-T Easy vector (Promega, Madison, WI). Resulting recombinant plasmids containing insert DNAs
5 within the expected size range were sequenced using T7 and Sp6 primers.

DNA sequencing

All the DNA sequencing reactions were performed using the BIGDYE™ Terminator Cycle sequencing kit (Perkin-Elmer, Wellesley, MA) with the sequences
10 being read on an automated ABI Prism 310 Genetic Analyzer (Applied Biosystems, Foster City, CA). Computer assessment of the DNA sequence information was performed using the MACVECTOR (Oxford Molecular, Madison, WI) software package.

cDNA library screening

The cDNA library was screened with digoxigenin labeled probes. A 258 bp DNA fragment amplified from the pGEM-deg6.4 clone using gene-specific forward (5'-GGCGGAGAATTTGTCCTGGAATGTCATTTGGTTTAG-3' (SEQ ID NO: 13)) and reverse (5'-GTACAATAGTGAGGTTGACAATG-3' (SEQ ID NO: 14)) primers; and a
20 374 bp DNA fragment amplified from the pBKS-CYPB3.843 clone with specific forward (5'-GGTGGTTGTGAATGCATG-3' (SEQ ID NO: 15)) and reverse (5'-TTATGCAGCAATAGGCTTGAAGACA-3' (SEQ ID NO: 16)) primers, were used to screen for CYP71Ds. The probes were labeled with digoxigenin-11-dUTP using the PCR DIG Labeling Mix (Roche Molecular Biochemicals, Indianapolis, IN), hybridized
25 to plaque lifts of the cDNA library plated at approximately 10,000 PFUs per 150 mm plate, and was hybridization detected with the DIG detection system according to the manufacturer's instructions (Roche Molecular Biochemicals, Indianapolis, IN). Plaques exhibiting strong hybridization were plaque purified, auto-subcloned to their plasmid forms according to the manufacturer's recommendations (Stratagene, La Jolla, CA), and
30 then subjected to DNA sequencing as described above.

RNA analysis

RNA gel blot analysis was carried out using 10 µg aliquots of total RNA. RNA samples were heat-denatured at 70°C for 15 minutes in sample buffer (1x MOPS, 50% formamide, 16% formaldehyde, 30 % glycerol, and 3% ethidium bromide), and size fractionated on a 1.2% agarose gel containing 1x MOPS and 18.1% formaldehyde. Uniformity of sample loading was determined by visual inspection of the gel for rRNA bands. The RNAs were then transferred to a Zeta Probe nylon membrane (Bio-Rad Laboratories, Hercules, CA) and hybridized according to the manufacturer's recommendations. Full-length cDNAs probes were labeled with [³²P]-dCTP (PRIME-IT Kit, Stratagene, La Jolla, CA) prior to hybridization. After hybridization, the membranes were washed in 2x SSC/0.1% SDS once at room temperature followed by sequential washes in 0.2x SSC/0.1% SDS at 42°C and 65°C. Hybridization was detected with a Phosphorimager (Molecular Dynamics, model 445 SI).

15 Construction of yeast expression vectors.

The coding regions of the P450 cDNAs were cloned into the pYeDP60 expression vector (Urban et al., J. Biol. Chem. 272:19176-19186, 1990; Pompon et al., Methods Enzymol. 272:51-64, 1996). Appropriate BamHI, EcoRI, and SstI restriction sites (underlined) were introduced via PCR primers containing these sequences either upstream of the translation start site (ATG) or downstream of the stop codon (TAA or TGA). The primers used to amplify the CYP71D20 cDNA were 5'-
 20 GGGGGATCCATGCAATTCCTCAGCTTGGTTTCC-3' (SEQ ID NO: 17) and 5'-
 GGGGAATTCTTACTCTCGAGAAGGTTGATAAGG-3' (SEQ ID NO: 18); for the
 CYP82E1 cDNA 5'- CCCGGATCCATGTATCATCTTCTTTCTCCC-3' (SEQ ID NO:
 25 19) and 5'- GGGGAATTCTCAATATTGATAAAGCGTAGGAGG-3' (SEQ ID NO:
 20) ; and for the CYP92A3 cDNA 5'-
 CCCGGATCCATGCAATCCTTCAGCTTGGTTTCC-3' (SEQ ID NO: 21) and 5'-
 GGGGAGCTCTCACTCGCAAGAAGATTGATAAGG-3' (SEQ ID NO: 22). Two
 long, overlapping (italicized) primers 5'-
 30 *GCCATTATCGGCGCAATACTAATCTCCAACTCCGCGGTAAAAAATTCAAGCT*
CCCACCTGGTCCAACAGCAGTC-3' (SEQ ID NO: 23) and 5'-
GGGGGATCCATGGACCTCCTCCTCATAGAAAAACCCTCGTCGCCTTATTG
CCGCCATTATCGGCGCAATACTA-3' (SEQ ID NO: 24) coding for the N-terminal

sequence of CYP73A1 (GenEMBL Z17369) up to the hinge region were used for the modification of the membrane anchoring segment of CYP73A27 to avoid possible problems with intracellular targeting due to the unusual N-terminus (Nedelkina et al., 1999); the reverse primer used for both amplifications was 5'-

5 GGGGAGCTCTTATGCAGCAATAGGCTTGAAGAC-3' (SEQ ID NO: 25).

CYP71D20 and CYP73A27 were amplified using full-length cDNA templates, whereas CYP82E1 and CYP92A5 were amplified directly from the cDNA library template. Amplifications were performed in 50 µl reactions containing 1x Pfx amplification buffer; 1 mM MgSO₄; 300 µM of each dNTP; 10 ng template DNA; 20 pmol each of

10 forward and reverse primer; and 1.25 units PLATINUM[®] Pfx Polymerase (Life Technologies, Rockville, MD). Reactions were preheated at 94°C for 2 minutes, followed by thirty-five cycles of denaturing at 94°C for 15 seconds, annealing at 55°C for 30 seconds, and elongating at 68° for 1.5 minutes. PCR products were ligated into the pGEM-T EASY vector (Promega, Madison, WI) and subcloned into the pYeDP60

15 vector. The resulting constructs were validated by a combination of PCR and DNA sequencing.

Yeast expression studies

Verified pYeDP60- P450 cDNA constructs were introduced into the yeast

20 WAT11 line, a derivative of the W303-1B strain (MAT a; ade 2-1; his 3-11; leu 2-3, -112; ura 3-1; can^R; cyr⁺), provided by Dr. P. Urban (Centre de Génétique Moléculaire, CNRS, Gif-sur-Yvette, France). The endogenous NADPH-cytochrome P450 reductase (CPR1) locus has been replaced with ATR1, a NADPH-cytochrome P450 reductase

25 from *Arabidopsis thaliana* (Pompon et al., Methods Enzymol. 272:51-64, 1996; Urban et al., J. Biol. Chem. 272:19176-19186, 1997), in the WAT11 line. Yeast was grown overnight in a 30°C shaker in YPAD (1g/l yeast extract; 1 g/l peptone; 20 g/l glucose; 200 mg/l adenine) liquid media. Cultures were harvested at an A₆₀₀ between 0.5 and 1.5. Cells were collected by centrifugation at 2,500 x g for 5 minutes at 4°C, and resuspended in ice-cold, sterile dH₂O. Cells were pelleted again as above and resuspended in 1 M

30 sorbitol. Forty µl of yeast suspension was mixed with 0.5 to 1 µg plasmid DNA (in <5 µl dH₂O) in a pre-chilled 0.5 ml tube, and transferred to a chilled cuvette with a 0.2 cm electrode gap. One pulse at 1.5 kV, 25 µF, and 200 Ohms was applied by an Eppendorf Electroporator (model 2510). A mixture of 500 µl of YPAD/1 M sorbitol was

immediately added to the electroporated cells. Cells were allowed to recover at 30°C for 1 hour, then spread onto SGI plates (1 g/l bactocasamino acids; 7 g/l yeast nitrogen base; 20 g/l glucose; 20 mg/l tryptophan; and 20 g/l agar). Transformed colonies appeared after 3 to 6 days of incubation at 30°C. Recombinant plasmids were confirmed by PCR
5 assays performed directly on randomly selected yeast colonies.

For expression studies, one colony was added to SGI media (1 g/l bactocasamino acids; 7 g/l yeast nitrogen base; 20 g/l glucose; and 20 mg/l tryptophan) and grown at 30°C for approximately 24 hours. An aliquot of this culture was diluted 1:50 into 250 ml of YPGE (10 g/l bactopectone; 10 g/l yeast extract; 5 g/l glucose; and 3% ethanol by
10 volume) and the cells were grown until all glucose was consumed. The absence of glucose was determined by placing a 200 µl aliquot of culture into a 1.5 ml tube, inserting a DIASTIX urinalysis reagent strip (Bayer, Elkhart, IN) for 30 seconds, and observing colorimetric changes indicating glucose levels. Induction was initiated by the addition of 5 grams of galactose (final concentration of 2%). The cultures were
15 maintained at 30°C for an additional 16 hours before collecting the cells by centrifugation at 7,000 x g for 10 minutes. The pelleted cells were washed with 100 ml of TES buffer (50 mM Tris-HCl pH, 7.5; 1 mM EDTA; 0.6 M sorbitol). The cells were centrifuged as above, resuspended in 100 ml of TES-M (TES supplemented with 10 mM 2-mercaptoethanol), and allowed to incubate at room temperature for 10 minutes. The
20 yeast cells were centrifuged again at 7,000 x g for 10 minutes, and the pellet was resuspended in 2.5 ml extraction buffer (1% bovine serum albumin, fraction V; 2 mM 2-mercaptoethanol; 1 mM phenylmethylsulfonyl fluoride, all dissolved in TES). Glass beads (0.5 mm in diameter, Biospec Products, Inc., Bartlesville, OK) were added until skimming the surface of the cell suspension. Cell walls were disrupted manually by hand
25 shaking in a cold room for 10 min at 30 second intervals separated by 30 second intervals on ice. Cell extracts were transferred to a 50 ml centrifuge tube, the glass beads were washed three times with 5 ml of extraction buffer, and the washes were pooled with the original cell extracts. Microsomes were prepared by differential centrifugation at 10,000 g for 10 minutes at 4°C to remove cellular debris, followed by
30 centrifugation at 100,000 x g for 70 minnutes at 4°C, and microsomal pellets were resuspended in 1.5 ml TEG-M buffer (50 mM Tris-HCl, pH 7.5; 1 mM EDTA; 20% glycerol; and 1.5 mM 2-mercaptoethanol) and stored frozen at -80°C until further assayed.

CO difference spectra

Fe²⁺ • CO vs. Fe²⁺ difference spectroscopy (Omura and Sato, J. Biol. Chem. 239:2370-2378, 1964) was performed using 0.4 ml of microsomes suspended in 1.6 ml of 50 mM Tris-HCl, pH 7.5; 1 mM EDTA; and 20% glycerol. A small amount of the reducing agent, sodium dithionite, was added, and the mixture was distributed between two cuvettes. A baseline was recorded between 400 and 500 nm on a Perkin Elmer Lambda 18 UV/visible spectrophotometer. CO was then bubbled into the sample cuvette for 1 minute, and the difference spectrum recorded again. The amount of functional P450 was estimated based on an absorbance coefficient of 91 mM⁻¹ • cm⁻¹.

5-epi-aristolochene-1, 3-hydroxylase assays

5-epi-aristolochene-1, 3-hydroxylase assays were performed in 0.5 ml polyethylene tubes in 100 µl volumes. 5-epi-aristolochene or 1-deoxycapsidiol dissolved in hexane was added to the tube, and the organic solvent was removed by incubation of the open tube at 30°C. 5-epi-aristolochene and 1-deoxycapsidiol were resuspended in 2 µl dimethyl sulfoxide before adding the reaction mixture. Reactions were carried out in 100 mM Tris-HCl, pH 7.5, to which microsomal protein was added to a final concentration of 1 mg/ml. Reactions were initiated by the addition of 2 mM NADPH. The final concentration of 5-epi-aristolochene and 1-deoxycapsidiol in these assays varied from 20 to 50 µM. After incubations for variable lengths of time at 30°C, the reactions were extracted with two volumes of ethyl acetate. The organic extracts were concentrated and evaluated by GC and GC-MS along with standards of 5-epi-aristolochene (Whitehead et al., Phytochemistry 28:775-779, 1989; Rising et al., J. Am. Chem. Soc. 122:1861-1866, 2000), 1-deoxycapsidiol (Whitehead et al., Phytochemistry 29:479-182, 1990), and capsidiol (Whitehead et al., Phytochemistry 26:1367-1369, 1987; Milat et al., Phytochemistry 30:2171-2173, 1991). GC analysis was routinely performed with an HP5890 GC equipped with a Hewlett-Packard HP-5 capillary column (30 m x 0.25 mm, 0.25 µm phase thickness) and FID as described previously (Rising et al., J. Am. Chem. Soc. 122:1861-1866, 2000). GC-MS analysis was performed at the University of Kentucky Mass Spectrometry Facility using a Varian 3400 gas chromatograph and a Finnigan INCOS 50 quadrupole mass selective detector. The GC was equipped with a J&W DB-5ms capillary column (15 m x 0.25 mm, 0.25 µm phase

thickness) and run with He as the carrier gas (10 psi.). Splitless injections were done at an injection port temperature of 280°C. The column temperature was maintained at 40°C for 1 minute and then increased to 280°C at 10°C per minute. Following separation by the GC column, samples were introduced directly into the electron impact
5 ionization source. Mass spectra were acquired at 70 eV, scanning from 40–440 Da in 1 second.

Production of cytochrome P450s

Using the standard molecular techniques described herein, the isolation of
10 additional cytochrome P450 coding sequences is readily accomplished. For example, using all or a portion of the amino acid sequence of any of the disclosed P450s, one may readily design P450-specific oligonucleotide probes, including P450 degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either
15 DNA strand and any appropriate portion of the P450 nucleotide sequence. General methods for designing and preparing such probes are provided, for example, in Ausubel et al., 2000, Current Protocols in Molecular Biology, Wiley Interscience, New York, and Berger and Kimmel, Guide to Molecular Cloning Techniques, 1987, Academic Press, New York. These oligonucleotides are useful for P450 gene isolation, either through
20 their use as probes capable of hybridizing to a P450 complementary sequence, or as primers for various amplification techniques, for example, polymerase chain reaction (PCR) cloning strategies.

Hybridization techniques and screening procedures are well known to those skilled in the art and are described, for example, in Ausubel et al. (supra); Berger and
25 Kimmel (supra); Chen et al., Arch. Biochem. Biophys. 324:255, 1995; and Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York. If desired, a combination of different oligonucleotide probes may be used for the screening of a recombinant DNA library. The oligonucleotides may be detectably-labeled using methods known in the art and used to probe filter replicas from
30 a recombinant DNA library. Recombinant DNA libraries are prepared according to methods well known in the art, for example, as described in Ausubel et al. (supra), or they may be obtained from commercial sources.

As discussed above, P450 oligonucleotides may also be used as primers in a polymerase chain reaction (PCR) amplification cloning strategy. PCR methods are well known in the art and are described, for example, in PCR Technology, Erlich, ed., Stockton Press, London, 1989; PCR Protocols: A Guide to Methods and Applications, Innis et al., eds., Academic Press, Inc., New York, 1990; and Ausubel et al. (supra). Primers are optionally designed to allow cloning of the amplified product into a suitable vector, for example, by including appropriate restriction sites at the 5' and 3' ends of the amplified fragment (as described herein). If desired, a P450 gene may be isolated using the PCR "RACE" technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al. (supra)). By this method, oligonucleotide primers based on a P450 sequence are oriented in the 3' and 5' directions and are used to generate overlapping PCR fragments. These overlapping 3'- and 5'-end RACE products are combined to produce an intact full-length cDNA. This method is described in Innis et al. (supra); and Frohman et al., Proc. Natl. Acad. Sci. USA 85:8998, (1988).

Additional methods for identifying sequences encoding P450s are provided in Maughan et al. (Arch. Biochem. Biophys. 341:104-111, 1997) and Clark et al. (Plant Mol. Biol. 33:875-885, 1997).

Useful P450 sequences may be isolated from any appropriate organism. Confirmation of a sequence's relatedness to a P450 polypeptide disclosed herein may be accomplished by a variety of conventional methods, for example, by comparing the sequence with a known p450 sequence found in a database. In addition, the activity of any P450 may be evaluated according to any of the techniques described herein.

P450 polypeptide expression

P450 polypeptides may be produced by transformation of a suitable host cell with all or part of a P450 DNA (for example, any one of the P450 cDNAs described herein) in a suitable expression vehicle or with a plasmid construct engineered for increasing the expression of a P450 polypeptide *in vivo*.

Those skilled in the field of molecular biology will appreciate that any of a wide variety of expression systems may be used to provide the recombinant protein. The precise host cell used is not critical to the invention. The P450 protein may be produced in a prokaryotic host, for example, *E. coli* TB1, or in a eukaryotic host, for example, *Saccharomyces cerevisiae*, insect cells, mammalian cells (for example, COS 1 or NIH

3T3 cells), or any of a number of plant cells including, without limitation, algae, tree species, ornamental species, temperate fruit species, tropical fruit species, vegetable species, legume species, monocots, dicots, or in any plant of commercial or agricultural significance. Particular examples of suitable plant hosts include, but are not limited to,

5 Conifers, Petunia, Tomato, Potato, Tobacco, Grape, Arabidopsis, Lettuce, Sunflower, Oilseed rape, Flax, Cotton, Sugarbeet, Celery, Soybean, Alfalfa, Medicago, Lotus, Vigna, Cucumber, Carrot, Eggplant, Cauliflower, Horseradish, Morning Glory, Poplar, Walnut, Apple, Asparagus, Grape, Rice, Maize, Millet, Onion, Barley, Orchard grass, Oat, Rye, Tobacco, and Wheat.

10 Such cells are available from a wide range of sources including: the American Type Culture Collection (Rockland, Md.); or from any of a number seed companies, for example, W. Atlee Burpee Seed Co. (Warminster, Pa.), Park Seed Co. (Greenwood, S.C.), Johnny Seed Co. (Albion, Me.), or Northrup King Seeds (Harstville, S.C.). Descriptions and sources of useful host cells are also found in Vasil I. K., Cell Culture

15 and Somatic Cell Genetics of Plants, Vol I, II, III Laboratory Procedures and Their Applications Academic Press, New York, 1984; Dixon, R. A., Plant Cell Culture--A Practical Approach, IRL Press, Oxford University, 1985; Green et al., Plant Tissue and Cell Culture, Academic Press, New York, 1987; and Gasser and Fraley, Science 244:1293, (1989).

20 For prokaryotic expression, DNA encoding a P450 polypeptide is carried on a vector operably linked to control signals capable of effecting expression in the prokaryotic host. If desired, the coding sequence may contain, at its 5' end, a sequence encoding any of the known signal sequences capable of effecting secretion of the expressed protein into the periplasmic space of the host cell, thereby facilitating

25 recovery of the protein and subsequent purification. Prokaryotes most frequently used are various strains of *E. coli*; however, other microbial strains may also be used. Plasmid vectors are used which contain replication origins, selectable markers, and control sequences derived from a species compatible with the microbial host. Examples of such vectors are found in Pouwels et al. (supra) or Ausubel et al. (supra). Commonly

30 used prokaryotic control sequences (also referred to as "regulatory elements") are defined herein to include promoters for transcription initiation, optionally with an operator, along with ribosome binding site sequences. Promoters commonly used to direct protein expression include the beta-lactamase (penicillinase), the lactose (lac), the

tryptophan (Trp) (Goeddel et al., Nucl. Acids Res. 8:4057 (1980)), and the tac promoter systems, as well as the lambda-derived P.sub.L promoter and N-gene ribosome binding site (Simatake et al., Nature 292:128 (1981)).

One particular bacterial expression system for P450 production is the *E. coli* pET expression system (Novagen). According to this expression system, DNA encoding a P450 is inserted into a pET vector in an orientation designed to allow expression. Since the P450 gene is under the control of the T7 regulatory signals, P450 expression is dependent on inducing the expression of T7 RNA polymerase in the host cell. This is typically achieved using host strains which express T7 RNA polymerase in response to IPTG induction. Once produced, recombinant P450 is then isolated according to standard methods known in the art, for example, those described herein.

Another bacterial expression system for P450 production is the pGEX expression system (Pharmacia). This system employs a GST gene fusion system that is designed for high-level expression of a gene or gene fragment as a fusion protein with rapid purification and recovery of the functional gene product. The P450 of interest is fused to the carboxyl terminus of the glutathione S-transferase protein from *Schistosoma japonicum* and is readily purified from bacterial lysates by affinity chromatography using Glutathione Sepharose 4B. Fusion proteins can be recovered under mild conditions by elution with glutathione. Cleavage of the glutathione S-transferase domain from the fusion protein is facilitated by the presence of recognition sites for site-specific proteases upstream of this domain. For example, proteins expressed in pGEX-2T plasmids may be cleaved with thrombin; those expressed in pGEX-3X may be cleaved with factor Xa.

Other prokaryotic systems useful for expressing eukaryotic P450s are described by Cooper (Mutat. Res. 454:45-52, 2000) and Dong et al. (Arch. Biochem. Biophys. 327:254-259, 1996). In addition, strategies for enhancing the prokaryotic expression of a cytochrome P450 in combination with cytochrome reductase are described in Porter et al. (Drug. Metab. Rev. 31:159-174, 1999).

For eukaryotic expression, the method of transformation or transfection and the choice of vehicle for expression of the P450 will depend on the host system selected. Transformation and transfection methods of numerous organisms, for example, the baker's yeast *Saccharomyces cerevisiae*, are described, e.g., in Ausubel et al. (supra); Weissbach and Weissbach, Methods for Plant Molecular Biology, Academic Press,

1989; Gelvin et al., Plant Molecular Biology Manual, Kluwer Academic Publishers, 1990; Kindle, K., Proc. Natl. Acad. Sci. U.S.A. 87:1228 (1990); Potrykus, I., Annu. Rev. Plant Physiol. Plant Mol. Biology 42:205 (1991); and BioRad (Hercules, Calif.) Technical Bulletin #1687 (Biolistic Particle Delivery Systems). Expression vehicles
5 may be chosen from those provided, e.g., in Cloning Vectors: A Laboratory Manual (P. H. Pouwels et al., 1985, Supp. 1987); Gasser and Fraley (supra); Clontech Molecular Biology Catalog (Catalog 1992/93 Tools for the Molecular Biologist, Palo Alto, Calif.); and the references cited above.

One preferred eukaryotic expression system is the mouse 3T3 fibroblast host cell
10 transfected with a pMAMneo expression vector (Clontech). pMAMneo provides: an RSV-LTR enhancer linked to a dexamethasone-inducible MMTV-LTR promoter, an SV40 origin of replication which allows replication in mammalian systems, a selectable neomycin gene, and SV40 splicing and polyadenylation sites. DNA encoding a P450 is inserted into the pMAMneo vector in an orientation designed to allow expression. The
15 recombinant P450 is then isolated as described below. Other preferable host cells which may be used in conjunction with the pMAMneo expression vehicle include COS cells and CHO cells (ATCC Accession Nos. CRL 1650 and CCL 61, respectively).

Alternatively, if desired, a P450 is produced by a stably-transfected mammalian cell line. A number of vectors suitable for stable transfection of mammalian cells are
20 available to the public, e.g., see Pouwels et al. (supra); methods for constructing such cell lines are also publicly available, e.g., in Ausubel et al. (supra). In one example, cDNA encoding the P450 is cloned into an expression vector which includes the dihydrofolate reductase (DHFR) gene. Integration of the plasmid and, therefore, the P450-encoding gene into the host cell chromosome is selected for by inclusion of 0.01-
25 300 μ M methotrexate in the cell culture medium (as described in Ausubel et al., supra). This dominant selection can be accomplished in most cell types. Recombinant protein expression can be increased by DHFR-mediated amplification of the transfected gene. Methods for selecting cell lines bearing gene amplifications are described in Ausubel et al. (supra); such methods generally involve extended culture in medium containing
30 gradually increasing levels of methotrexate. DHFR-containing expression vectors commonly used for this purpose include pCVSEII-DHrF and pAdD26SV(A) (described in Ausubel et al., supra). Any of the host cells described above or, preferably, a DHFR-deficient CHO cell line (for example, CHO DHFR cells, ATCC Accession Number CRL

9096) are among the host cells preferred for DHFR selection of a stably-transfected cell line or DHFR-mediated gene amplification.

A cytochrome P450 may also be produced in insect cells, such cells include, without limitation, *Spodoptera frugiperda* (Sf)-9, Sf-21, or *Drosophila melanogaster* Schneider (SL-2) cells. For P450 production, insect cells are typically infected with a baculovirus, for example, *Autographa californica* Multiple Nuclear Polyhedrosis Virus (AcMNPV) containing an expression cassette for such a protein, e.g., cytochrome P450, at a multiplicity of infection of 1 to 10. The infected cells are generally cultured in a standard insect cell culture medium for 24 to 48 hours prior to recovering the protein using standard molecular biology techniques. If desired, a P450 polypeptide may also be produced in insect cells directly transfected with a DNA construct containing an expression cassette encoding the P450.

Furthermore, any of the cytochrome P450s described herein may be produced in yeast, for example, *Pichia pastoris*. In order to produce the P450, yeast cells are transformed with an expression cassette containing, for example, a promoter such as the AOX1 or phosphoglycerate kinase gene promoter, the P450 gene to be expressed, and a terminator. Such an expression cassette may contain an origin of replication or it may be integrated into the yeast genomic DNA. The expression cassette is generally introduced by lithium acetate transformation or by the use of spheroplasts. In order to select for successfully transformed cells, the yeast are plated, for example, on minimal media which only allows yeast carrying the introduced expression cassette to grow.

In addition, expression of recombinant proteins in yeast using a *Hansenula polymorpha* expression system is described in U.S. Patent Nos. 5,741,674 and 5,672,487.

A P450 may also be produced by a stably-transfected plant cell line or by a transgenic plant. Such genetically-engineered plants are useful for a variety of industrial and agricultural applications as discussed below. Importantly, this invention is applicable to gymnosperms and angiosperms, and will be readily applicable to any new or improved transformation or regeneration method.

A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants are available to the public; such vectors are described in Pouwels et al. (supra), Weissbach and Weissbach (supra), and Gelvin et al. (supra). Methods for constructing such cell lines are described in, e.g., Weissbach and

Weissbach (supra), and Gelvin et al. (supra). Typically, plant expression vectors include (1) a cloned P450 gene under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (for example, one conferring inducible or
5 constitutive expression, or environmentally- or developmentally-regulated, or pathogen- or wound-inducible, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

The P450 DNA sequence of the invention may, if desired, be combined with other
10 DNA sequences in a variety of ways. The P450 DNA sequence of the invention may be employed with all or part of the gene sequences normally associated with a P450. In its component parts, a DNA sequence encoding a P450 is combined in a DNA construct having a transcription initiation control region capable of promoting transcription and translation in a host cell.

15 In general, the constructs will involve regulatory regions functional in plants which provide for production of a P450 as discussed herein. The open reading frame coding for the P450, or a functional fragment thereof, will be joined at its 5' end to a transcription initiation regulatory region such as the sequence naturally found in the 5' upstream region of a P450 structural gene, for example, a CYP71D20 (SEQ ID NO:2) or
20 CYP71D21 (SEQ ID NO:4) gene. Numerous other transcription initiation regions are available which provide for constitutive or inducible regulation.

For applications when developmental, cell, tissue, hormonal, environmental, or pathogen-inducible expression are desired, appropriate 5' upstream non-coding regions are obtained from other genes; for example, from genes regulated during seed
25 development, embryo development, leaf development, or in response to a pathogen.

Regulatory transcript termination regions may also be provided in DNA constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding a P450 or any convenient transcription termination region derived from a different gene source. The transcript termination region will contain preferably
30 at least 1-3 kb of sequence 3' to the structural gene from which the termination region is derived.

An example of a useful plant promoter according to the invention is a caulimovirus promoter, such as, a cauliflower mosaic virus (CaMV) promoter. These promoters confer high levels of expression in most plant tissues, and the activity of these promoters is not dependent on virally encoded proteins. CaMV is a source for both the 35S and 19S promoters. In most tissues of transgenic plants, the CaMV 35S promoter is a strong promoter (see, e.g., Odell et al., *Nature* 313:810 (1985)). The CaMV promoter is also highly active in monocots (see, e.g., Dekeyser et al., *Plant Cell* 2:591 (1990); Terada and Shimamoto, *Mol. Gen. Genet.* 220:389, (1990)). Moreover, activity of this promoter can be further increased (i.e., between 2-10 fold) by duplication of the CaMV 35S promoter (see e.g., Kay et al., *Science* 236:1299 (1987); Ow et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:4870 (1987); and Fang et al., *Plant Cell* 1:141 (1989)). Other useful plant promoters include, without limitation, the nopaline synthase promoter (An et al., *Plant Physiol.* 88:547 (1988)) and the octopine synthase promoter (Fromm et al., *Plant Cell* 1:977 (1989)).

For certain applications, it may be desirable to produce the P450 gene product in an appropriate tissue, at an appropriate level, or at an appropriate developmental time. For this purpose, there is an assortment of gene promoters, each with its own distinct characteristics embodied in its regulatory sequences, which have been shown to be regulated in response to the environment, hormones, and/or developmental cues. These include gene promoters that are responsible for heat-regulated gene expression (see, e.g., Callis et al., *Plant Physiol.* 88:965 (1988); Takahashi and Komeda, *Mol. Gen. Genet.* 219:365 (1989); and Takahashi et al., *Plant J.* 2:751(1992)), light-regulated gene expression (e.g., the pea *rbcS*-3A described by Kuhlemeier et al. (*Plant Cell* 1:471 (1989); the maize *rbcS* promoter described by Schaffner and Sheen, (*Plant Cell* 3:997 (1991); or the chlorophyll a/b-binding protein gene found in pea described by Simpson et al. (*EMBO J.* 4:2723 (1985)), hormone-regulated gene expression (for example, the abscisic acid (ABA) responsive sequences from the *Em* gene of wheat described by Marcotte et al. (*Plant Cell* 1:969 (1989); the ABA-inducible HVA1 and HVA22, and the *rd29A* promoters described for barley and *Arabidopsis* by Straub et al. (*Plant Cell* 6:617 (1994), Shen et al. (*Plant Cell* 7:295 (1994)), and wound-induced gene expression (for example, of *wun1* described by Siebertz et al. (*Plant Cell* 1:961 (1989)), or organ-specific gene expression (for example, of the tuber-specific storage protein gene described by Roshal et al. (*EMBO J.* 6:1155 (1987); the 23-kDa *zein* gene from maize

described by Schernthaner et al. (EMBO J. 7:1249 (1988); or the French bean beta-phaseolin gene described by Bustos et al., (Plant Cell 1:839 (1989)); and pathogen-inducible gene expression described by Chappell et al. in U.S. Ser. Nos. 08/471,983, 08/443,639, and 08/577,483, hereby incorporated by reference.

5 Plant expression vectors may also optionally include RNA processing signals, for example, introns, which have been shown to be important for efficient RNA synthesis and accumulation (Callis et al., Genes and Dev. 1:1183 (1987)). The location of the RNA splice sequences can dramatically influence the level of transgene expression in plants. In view of this fact, an intron may be positioned upstream or downstream of a
10 P450-encoding sequence in the transgene to modulate levels of gene expression.

In addition to the aforementioned 5' regulatory control sequences, the expression vectors may also include regulatory control regions which are generally present in the 3' regions of plant genes (Thornburg et al., Proc. Natl. Acad. Sci. U.S.A. 84:744 (1987); An et al., Plant Cell 1:115 (1989)). For example, the 3' terminator region may be
15 included in the expression vector to increase stability of the mRNA. One such terminator region may be derived from the PI-II terminator region of potato. In addition, other commonly used terminators are derived from the octopine or nopaline synthase signals.

The plant expression vector also typically contains a dominant selectable marker
20 gene used to identify those cells that have become transformed. Useful selectable genes for plant systems include genes encoding antibiotic resistance genes, for example, those encoding resistance to hygromycin, kanamycin, bleomycin, G418, streptomycin, or spectinomycin. Genes required for photosynthesis may also be used as selectable markers in photosynthetic-deficient strains. Alternatively, the green-fluorescent protein
25 from the jellyfish *Aequorea victoria* may be used as a selectable marker (Sheen et al., Plant J. 8:777, 1995; Chiu et al., Current Biology 6:325 (1996)). Finally, genes encoding herbicide resistance may be used as selectable markers; useful herbicide resistance genes include the *bar* gene encoding the enzyme phosphinothricin acetyltransferase and conferring resistance to the broad-spectrum herbicide BASTA
30 (Hoechst AG, Frankfurt, Germany).

Efficient use of selectable markers is facilitated by a determination of the susceptibility of a plant cell to a particular selectable agent and a determination of the concentration of this agent which effectively kills most, if not all, of the transformed

cells. Some useful concentrations of antibiotics for tobacco transformation include, e.g., 75-100 µg/ml (kanamycin), 20-50 µg/ml (hygromycin), or 5-10 µg/ml (bleomycin). A useful strategy for selection of transformants for herbicide resistance is described, e.g., by Vasil et al., *supra*.

- 5 It should be readily apparent to one skilled in the art of molecular biology, especially in the field of plant molecular biology, that the level of gene expression is dependent, not only on the combination of promoters, RNA processing signals, and terminator elements, but also on how these elements are used to increase the levels of selectable marker gene expression.

10

Plant transformation

- Upon construction of the plant expression vector, several standard methods are available for introduction of the vector into a plant host, thereby generating a transgenic plant. These methods include (1) *Agrobacterium*-mediated transformation (*A. tumefaciens* or *A. rhizogenes*) (see, e.g., Lichtenstein and Fuller, In: Genetic Engineering, vol. 6, PWJ Rigby, ed, London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: DNA Cloning, Vol II, D. M. Glover, ed, Oxford, IRI Press, 1985), (2) the particle delivery system (see, e.g., Gordon-Kamm et al., *Plant Cell* 2:603 (1990); or BioRad Technical Bulletin 1687, *supra*), (3) microinjection protocols (see, 15 e.g., Green et al., *supra*), (4) polyethylene glycol (PEG) procedures (see, e.g., Draper et al., *Plant Cell Physiol.* 23:451 (1982); or e.g., Zhang and Wu, *Theor. Appl. Genet.* 76:835 (1988)), (5) liposome-mediated DNA uptake (see, e.g., Freeman et al., *Plant Cell Physiol.* 25:1353 (1984)), (6) electroporation protocols (see, e.g., Gelvin et al., *supra*; Dekeyser et al., *supra*; Fromm et al., *Nature* 319:791 (1986); Sheen, *Plant Cell* 2:1027 20 (1990); or Jang and Sheen, *Plant Cell* 6:1665 (1994)), and (7) the vortexing method (see, e.g., Kindle, *supra*). The method of transformation is not critical to the present invention. Any method which provides for efficient transformation may be employed. As newer methods are available to transform crops or other host cells, they may be directly applied.

- 30 The following is an example outlining one particular technique, an *Agrobacterium*-mediated plant transformation. By this technique, the general process for manipulating genes to be transferred into the genome of plant cells is carried out in two phases. First, cloning and DNA modification steps are carried out in *E. coli*, and the

plasmid containing the gene construct of interest is transferred by conjugation or electroporation into *Agrobacterium*. Second, the resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent introduction into plants. Resistance genes can be carried on the vector, one for selection in bacteria, for example, streptomycin, and another that will function in plants, for example, a gene encoding kanamycin resistance or herbicide resistance. Also present on the vector are restriction endonuclease sites for the addition of one or more transgenes and directional T-DNA border sequences which, when recognized by the transfer functions of *Agrobacterium*, delimit the DNA region that will be transferred to the plant.

In another example, plant cells may be transformed by shooting into the cell tungsten microprojectiles on which cloned DNA is precipitated. In the Biolistic Apparatus (Bio-Rad) used for the shooting, a gunpowder charge (22 caliber Power Piston Tool Charge) or an air-driven blast drives a plastic macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to pass through. As a result, the plastic macroprojectile smashes against the stopping plate, and the tungsten microprojectiles continue toward their target through the hole in the plate. For the present invention, the target can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

In general, transfer and expression of transgenes in plant cells are now routine practices to those skilled in the art, and have become major tools to carry out gene expression studies in plants and to produce improved plant varieties of agricultural or commercial interest.

Transgenic plant regeneration

Plants cells transformed with plant expression vectors can be regenerated, for example, from single cells, callus tissue, or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from

almost any plant can be successfully cultured to regenerate an entire plant; such techniques are described, e.g., in Vasil *supra*; Green et al., *supra*; Weissbach and Weissbach, *supra*; and Gelvin et al., *supra*.

In one particular example, a cloned P450, under the control of the EAS4 promoter and the nopaline synthase terminator and carrying a selectable marker (for example, kanamycin resistance), is transformed into *Agrobacterium*. Transformation of leaf discs (for example, of tobacco leaf discs), with vector-containing *Agrobacterium* is carried out as described by Horsch et al. (*Science* 227:1229 (1985)). Putative transformants are selected after a few weeks (for example, 3 to 5 weeks) on plant tissue culture media containing kanamycin (e.g., 100 µg/ml). Kanamycin-resistant shoots are then placed on plant tissue culture media without hormones for root initiation. Kanamycin-resistant plants are then selected for greenhouse growth. If desired, seeds from self-fertilized transgenic plants can then be sowed in soil-less medium and grown in a greenhouse. Kanamycin-resistant progeny are selected by sowing surface sterilized seeds on hormone-free kanamycin-containing media. Analysis for the integration of the transgene is accomplished by standard techniques (see, for example, Ausubel et al., *supra*; Gelvin et al., *supra*).

Transgenic plants expressing the selectable marker are then screened for transmission of the transgene DNA by standard immunoblot and DNA detection techniques. Each positive transgenic plant and its transgenic progeny is unique in comparison to other transgenic plants established with the same transgene. Integration of the transgene DNA into the plant genomic DNA is in most cases random, and the site of integration can profoundly affect the levels and the tissue and developmental patterns of transgene expression. Consequently, a number of transgenic lines are usually screened for each transgene to identify and select plants with the most appropriate expression profiles.

Transgenic lines are generally evaluated for levels of transgene expression. Expression at the RNA level is determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis are employed and include PCR amplification assays using oligonucleotide primers designed to amplify only transgene RNA templates and solution hybridization assays using transgene-specific probes (see, e.g., Ausubel et al., *supra*). The RNA-positive plants are then analyzed for protein expression by Western immunoblot analysis using specific antibodies to the P450 (see,

e.g., Ausubel et al., supra). In addition, *in situ* hybridization and immunocytochemistry according to standard protocols can be done using transgene-specific nucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue.

Once the recombinant P450 is expressed in any cell or in a transgenic plant (for example, as described above), it may be isolated, e.g., using affinity chromatography. In one example, an anti-P450 antibody (e.g., produced as described in Ausubel et al., supra, or by any standard technique) may be attached to a column and used to isolate the polypeptide. Lysis and fractionation of P450-producing cells prior to affinity chromatography may be performed by standard methods (see, e.g., Ausubel et al., supra). Once isolated, the recombinant protein can, if desired, be further purified, for example, by high performance liquid chromatography (see, e.g., Fisher, Laboratory Techniques in Biochemistry and Molecular Biology, eds., Work and Burdon, Elsevier, 1980).

These general techniques of polypeptide expression and purification can also be used to produce and isolate useful P450 fragments or analogs.

Use

The aforementioned cytochrome P450 polypeptides of the invention are useful in the biosynthesis of hormones, lipids, and secondary metabolites, and may also help plants tolerate potentially harmful exogenous chemicals such as herbicides, pesticides, and pollutants. In addition, such cytochrome P450 polypeptides are useful in the chemical defense of plants against insects, as well as against bacterial, viral, and fungal infection.

Engineering plant disease resistance

Plasmid constructs designed for the expression of a P450 gene product are useful, for example, for activating plant defense pathways that confer anti-pathogenic properties to a transgenic plant, for example, the production of phytoalexins. P450 genes that are isolated from a host plant (e.g., *Nicotiana*) may be engineered for expression in the same plant, a closely related species, or a distantly related plant species. For example, a P450 gene may be engineered for constitutive low-level expression and then transformed into a *Nicotiana* host plant. Alternatively, the P450 gene may be engineered for expression in other solanaceous plants, including, but not

limited to, potato and tomato. To achieve pathogen resistance, it is important to express a P450 protein at an effective level. Evaluation of the level of pathogen protection conferred to a plant by ectopic expression of the P450 gene is determined according to conventional methods and assays.

5

Industrial applications

The invention also includes engineering host cells to include novel isoprenoid metabolic pathways useful in the production of new isoprenoid compounds. By introducing genes encoding an isoprenoid synthase (as disclosed in U.S. Patent No. 10 5,824,774 and WO 00/17327) and a cytochrome P450, an acetyltransferase, a methyl transferase, a fatty acyltransferase, or a combination thereof, various isoprenoid reaction products may be modified, controlled, or manipulated, resulting in enhancement of production of numerous isoprenoid reaction products, for example, the production of novel monoterpenes, diterpenes, and sesquiterpenes. Such compounds are useful as 15 phytoalexins, insecticides, perfumes, and pharmaceuticals such as anti-bacterial and fungal agents.

In one working example, an isoprenoid synthase or a chimeric isoprenoid synthase (as disclosed in U.S. Patent 5,824,774 and WO 00/17327) and a P450 gene are introduced into yeast, for example, using any of the procedures described herein. If 20 desired, such cells may also express, either independently or in combination, an acetyltransferase (see, for example, Walker et al., Proc. Natl. Acad. Sci. USA 18:583-587, 2000), a methylase transferase (see, for example, Diener et al., Plant Cell 12:853-870, 2000) gene, or a fatty acyltransferase gene, as well as a cytochrome reductase. Cells are then cultured under standard conditions and the production of isoprenoid 25 compounds is assayed according to methods known in the art. Isoprenoid compounds are further purified according to methods well known in the art. Cells expressing novel isoprenoid compounds are taken as useful in the invention.

Such methods provide a unique approach for producing novel isoprenoid starting materials and end products. Either prokaryotic or eukaryotic cells transformed with any 30 of the aforementioned enzymes (or combinations thereof) may be used. Moreover, isoprenoid compounds may be produced in any number of ways known in the art including an *in vitro* combination of purified enzymes with an appropriate substrate or direct fermentation using a host cell which expresses any combination of the

aforementioned enzymes and the appropriate substrates sufficient to drive production of isoprenoid compounds.

The invention is also useful for the production of insect attractants and deterrents, which may either deter insect pests or attract insect predators. In addition,
5 the invention is also useful for generating novel flavorings and perfumes.

Other embodiments

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and can make various changes and
10 modifications of the invention to adapt it to various usages and conditions. Thus, other embodiments are also within the claims.

All publications and patents mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent was specifically and individually indicated to be incorporated by reference.

15

We claim:

CLAIMS

1. A method for producing an isoprenoid compound, said method comprising the steps of:
- 5 (a) culturing a host cell under conditions suitable for expressing a first recombinant protein comprising an isoprenoid synthase and a second recombinant protein comprising a cytochrome P450 polypeptide, wherein said first and second recombinant protein catalyze the formation of an isoprenoid compound not normally produced by said host cell; and
- 10 (b) recovering said isoprenoid compound.
2. The method of claim 1, wherein said cytochrome P450 polypeptide is selected from the group consisting of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, a polypeptide comprising the amino acid sequence of SEQ ID NO:3, a
- 15 polypeptide comprising the amino acid sequence of SEQ ID NO:5, and a polypeptide comprising the amino acid sequence of SEQ ID NO:7.
3. The method of claim 1, wherein said host cell further comprises a third recombinant protein selected from the group consisting of an acetyltransferase, a
- 20 methyltransferase, and a fatty acyltransferase.
4. The method of claim 1, wherein said host cell expresses an endogenous or recombinant cytochrome reductase.
- 25 5. The method of claim 1, wherein said host cell is a yeast cell, a bacterial cell, an insect cell, or a plant cell.
6. A host cell expressing a recombinant isoprenoid synthase and a recombinant cytochrome P450 polypeptide.
- 30 7. The host cell of claim 6, wherein said host cell further expresses a recombinant acetyltransferase, a recombinant methyltransferase, or a recombinant fatty acyltransferase.

8. The host cell of claim 6, wherein said host cell expresses an endogenous or recombinant cytochrome reductase.

5 9. The host cell of claim 6, wherein said host cell is selected from the group consisting of a yeast cell, a bacterial cell, an insect cell, and a plant cell.

10 10. An isolated cytochrome P450 polypeptide having 80% identity to the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:3.

11. The polypeptide of claim 10, said polypeptide comprising the amino acid sequence of SEQ ID NO:1.

15 12. The polypeptide of claim 10, said polypeptide comprising the amino acid sequence of SEQ ID NO:3.

13. An isolated cytochrome P450 polypeptide having 85% identity to the amino acid sequence of SEQ ID NO:5 or SEQ ID NO:7.

20 14. The polypeptide of claim 13, said polypeptide comprising the amino acid sequence of SEQ ID NO:5.

25 15. The polypeptide of claim 13, said polypeptide comprising the amino acid sequence of SEQ ID NO:7.

16. An isolated cytochrome P450 polypeptide having 97% identity to the amino acid sequence of SEQ ID NO:11.

30 17. The polypeptide of claim 16, said polypeptide comprising the amino acid sequence of SEQ ID NO:11.

18. An isolated cytochrome P450 nucleic acid molecule which encodes a polypeptide having 80% identity to the amino acid sequence of SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:11.
- 5 19. The isolated nucleic acid molecule of claim 18, wherein said nucleic acid molecule comprises SEQ ID NO:1.
20. The isolated nucleic acid molecule of claim 18, wherein said nucleic acid molecule
- 10 comprises SEQ ID NO:3.
21. The isolated nucleic acid molecule of claim 18, wherein said nucleic acid molecule comprises SEQ ID NO:11.
- 15 22. An isolated cytochrome P450 nucleic acid molecule which encodes a polypeptide having 85% identity to the amino acid sequence of SEQ ID NO:5 or SEQ ID NO:7.
23. The isolated nucleic acid molecule of claim 22, wherein said nucleic acid
- 20 molecule comprises SEQ ID NO:5.
24. The isolated nucleic acid molecule of claim 22, wherein said nucleic acid molecule comprises SEQ ID NO:7.
- 25 25. An isolated nucleic acid molecule that specifically hybridizes under highly stringent conditions to any one of the complement of the sequence described in SEQ ID NOS:2, 4, 6, 8, or 12, wherein said nucleic acid encodes a cytochrome P450 polypeptide.
- 30 26. A vector comprising an isolated nucleic acid molecule selected from the group consisting of the isolated nucleic acid molecule of any one of claims 18-21 and 22-24.

27. A host cell comprising an isolated nucleic acid molecule selected from the group consisting of the isolated nucleic acid molecule of any one of claims 18-21 and 22-24.

5 28. A plant or plant component comprising an isolated nucleic acid molecule selected from the group consisting of the isolated nucleic acid molecule of any one of claims 18-21 and 22-24.

10 29. A process for conferring disease resistance on a plant or plant component, said method comprising the steps of:

(a) introducing into a plant cell a transgene comprising a nucleic acid molecule selected from the group consisting of the isolated nucleic acid molecule of any one of claims 18-21 and 22-24 operably linked to a promoter functional in said plant cell to yield a transformed plant cell; and

15 (b) regenerating a plant or plant component from said transformed plant cell, wherein the nucleic acid molecule is expressed in said plant or plant component, thereby conferring disease resistance on said plant or plant component.

20 30. A method of producing a cytochrome P450 polypeptide in a host cell, said method comprising the steps of:

(a) providing a host cell of claim 27;

(b) culturing said host cell under conditions for expressing said isolated nucleic acid molecule encoding said cytochrome P450 polypeptide; and

(c) recovering said cytochrome P450 enzyme.

25

31. A method for producing an altered compound, said method comprising the step of contacting said compound with the isolated polypeptide of any one of claims 10-15 under conditions allowing for the hydroxylation, oxidation, demethylation, methylation, or any combination of said enzymatic activity of said compound and
30 recovering said altered compound.

32. A hydroxylating agent comprising the isolated polypeptide of any one of claims 10-15.

1/8




Figure 1

2/8




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

7/8




Figure 7




Figure 8

SEQUENCE LISTING

<110> University of Kentucky Research Foundation

<120> Cytochrome P450s and Uses Thereof

<130> 07678/100W03

<150> US 60/274,241

<151> 2001-03-09

<150> US 60/275,597

<151> 2001-03-13

<160> 43

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 473

<212> PRT

<213> Nicotiana tabacum

<400> 1

```

Met Gln Phe Phe Ser Leu Val Ser Ile Phe Leu Phe Leu Ala Phe Leu
 1           5           10           15
Phe Leu Leu Arg Lys Trp Lys Asn Ser Asn Ser Gln Ser Lys Lys Leu
      20           25           30
Pro Pro Gly Pro Trp Lys Ile Pro Ile Leu Gly Ser Met Leu His Met
      35           40           45
Ile Gly Gly Glu Pro His His Val Leu Arg Asp Leu Ala Lys Lys Tyr
      50           55           60
Gly Pro Leu Met His Leu Gln Leu Gly Glu Ile Ser Ala Val Val Val
65           70           75           80
Thr Ser Arg Asp Met Ala Lys Glu Val Leu Lys Thr His Asp Val Val
      85           90           95
Phe Ala Ser Arg Pro Lys Ile Val Ala Met Asp Ile Ile Cys Tyr Asn
      100          105          110
Gln Ser Asp Ile Ala Phe Ser Pro Tyr Gly Asp His Trp Arg Gln Met
      115          120          125
Arg Lys Ile Cys Val Met Glu Leu Leu Asn Ala Lys Asn Val Arg Ser
      130          135          140
Phe Ser Ser Ile Arg Arg Asp Glu Val Val Arg Leu Ile Asp Ser Ile
145          150          155          160
Arg Ser Asp Ser Ser Gly Glu Leu Val Asn Phe Thr Gln Arg Ile
      165          170          175
Ile Trp Phe Ala Ser Ser Met Thr Cys Arg Ser Ala Phe Gly Gln Val
      180          185          190
Leu Lys Gly Gln Asp Ile Phe Ala Lys Lys Ile Arg Glu Val Ile Gly

```

195	200	205
Leu Ala Glu Gly Phe Asp Val Val Asp Ile Phe Pro Thr Tyr Lys Phe		
210	215	220
Leu His Val Leu Ser Gly Met Lys Arg Lys Leu Leu Asn Ala His Leu		
225	230	235
Lys Val Asp Ala Ile Val Glu Asp Val Ile Asn Glu His Lys Lys Asn		240
	245	250
Leu Ala Ala Gly Lys Ser Asn Gly Ala Leu Glu Asp Met Phe Ala Ala		255
	260	265
Gly Thr Glu Thr Ser Ser Thr Thr Thr Val Trp Ala Met Ala Glu Met		270
	275	280
Met Lys Asn Pro Ser Val Phe Thr Lys Ala Gln Ala Glu Val Arg Glu		285
	290	295
Ala Phe Arg Asp Lys Val Ser Phe Asp Glu Asn Asp Val Glu Glu Leu		300
305	310	315
Lys Tyr Leu Lys Leu Val Ile Lys Glu Thr Leu Arg Leu His Pro Pro		320
	325	330
Ser Pro Leu Leu Val Pro Arg Glu Cys Arg Glu Asp Thr Asp Ile Asn		335
	340	345
Gly Tyr Thr Ile Pro Ala Lys Thr Lys Val Met Val Asn Val Trp Ala		350
	355	360
Leu Gly Arg Asp Pro Lys Tyr Trp Asp Asp Ala Glu Ser Phe Lys Pro		365
	370	375
Glu Arg Phe Glu Gln Cys Ser Val Asp Phe Phe Gly Asn Asn Phe Glu		380
385	390	395
Phe Leu Pro Phe Gly Gly Gly Arg Arg Ile Cys Pro Gly Met Ser Phe		400
	405	410
Gly Leu Ala Asn Leu Tyr Leu Pro Leu Ala Gln Leu Leu Tyr His Phe		415
	420	425
Asp Trp Lys Leu Pro Thr Gly Ile Met Pro Arg Asp Leu Asp Leu Thr		430
	435	440
Glu Leu Ser Gly Ile Thr Ile Ala Arg Lys Gly Asp Leu Tyr Leu Asn		445
	450	455
Ala Thr Pro Tyr Gln Pro Ser Arg Glu		460
465	470	

<210> 2

<211> 1660

<212> DNA

<213> Nicotiana tabacum

<400> 2

```

ggatgggtcta ataatcctcc atttatctcc gaaaatgcaa ttcttcagct tggtttccat 60
tttcctcttc ctagctttcc tatttttgtt gaggaatgg aagaactcca atagccaaag 120
caaaaaattg ccaccaggtc catggaaaat accaatacta ggaagtatgc ttcatatgat 180
tggtggagaa cgcaccatg tccttagaga tttagccaaa aaatatggac cacttatgca 240
ccttcagtta ggtgaaattt ctgcagttgt ggttacttct agggacatgg caaaagaagt 300
gctaaaaact catgacgtcg tttttgcata taggcctaaa attgtagcca tggacattat 360
ctgttataac cagtccgaca ttgcctttag cccttatggc gaccactgga gacaaatgca 420
taaaatttgt gtcatggaac ttctcaatgc aaagaatggt cggtctttca gctccatcag 480

```

```

acgtgatgaa gtcgttcgtc tcattgactc tatccgggtca gattcttctt caggtgagct 540
agttaatttt acgcagagga tcatttggtt tgcaagctcc atgacgtgta gatcagcatt 600
tgggcaagta ctcaaggggc aagacatatt tgccaaaaag atcagagaag taataggatt 660
agcagaaggc tttgatgtgg tagacatctt ccctacatac aagtttcttc atgttctcag 720
tgggatgaag cgtaaacttt tgaatgccca ccttaaggta gacgccattg ttgaggatgt 780
catcaacgag cacaagaaaa atcttgacgc tggcaaaagt aatggcgcat tagaggacat 840
gtttgctgcc ggaacagaaa cttcatcaac aacaactgta tgggctatgg ctgaaatgat 900
gaagaatcca agtgatttca ccaaagctca agcagaagtg cgagaagcct ttagggacaa 960
agtatctttt gatgaaaatg atgtggagga gctgaaatac ttaaagttag tcattaaaga 1020
aactttgaga cttcatccac cgtctccact tttgggtcca agagaatgca gggaagatac 1080
ggatataaac ggctacacta ttcttgcaaa gaccaaagtt atgggtaatg ttggggcatt 1140
gggaagagat ccaaaatatt gggatgacgc ggaaagcttt aagccagaga gatttgagca 1200
atgctctgtg gatttttttg gtaataattt tgagtttctt ccctttggcg gtggacggag 1260
aatgtgtcct ggaatgtcat ttggtttagc taatctttac ttgccattgg ctcaattact 1320
ctatcacttt gactggaaac tcccaaccgg aatcatgcca agagacttag acttgaccga 1380
attatcgggg ataactattg ctagaaaggg tgacctttac ttaaagcca ctccttatca 1440
accttctcga gagtaattca atattggcat aaacatttta aatttccttc atcaacctca 1500
atattgtaca ataatcattc ttctgggtgt ataggcttta tcgatttcca atacatgtat 1560
tctttattaa aaaatgtatc acattccatg tagaaggagg acgcaccaat taattgtgcc 1620
atgattttag ggtaacttgt tccatcttaa aaaaaaaaaa 1660

```

<210> 3

<211> 474

<212> PRT

<213> *Nicotiana tabacum*

<400> 3

```

Met Glu Phe Phe Ser Leu Val Ser Ile Phe Leu Phe Leu Ser Phe Leu
 1           5           10           15
Phe Leu Leu Arg Lys Cys Lys Asn Ser Asn Ser Gln Thr Lys Gln Leu
 20           25           30
Pro Pro Gly Pro Trp Lys Ile Pro Ile Leu Gly Ser Met Leu His Met
 35           40           45
Leu Gly Gly Glu Pro His His Ile Leu Arg Asp Leu Ala Lys Lys Tyr
 50           55           60
Gly Pro Ile Met His Leu Gln Phe Gly Glu Ile Ser Ala Val Val Val
 65           70           75           80
Thr Ser Arg Glu Met Ala Lys Glu Val Leu Lys Thr His Asp Val Val
 85           90           95
Phe Ala Ser Arg Pro Lys Ile Val Ala Met Asp Ile Ile Cys Tyr Asn
100           105           110
Gln Ser Asp Ile Ala Phe Ser Pro Tyr Gly Asp His Trp Arg Gln Met
115           120           125
Arg Lys Ile Cys Val Met Glu Leu Leu Asn Ala Lys Asn Val Arg Ser
130           135           140
Phe Ser Ser Ile Arg Arg Asp Glu Val Val Arg Leu Ile Asp Ser Ile
145           150           155           160
Arg Ser Asp Ser Ser Gly Glu Leu Val Asn Phe Thr Gln Arg Ile
165           170           175
Ile Trp Phe Ala Ser Ser Met Thr Cys Arg Ser Ala Phe Gly Gln Val
180           185           190

```

Leu Lys Gly Gln Asp Val Phe Ala Lys Lys Ile Arg Glu Val Ile Gly
 195 200 205
 Leu Ala Glu Gly Phe Asp Val Ala Asp Ile Phe Pro Ser Tyr Lys Phe
 210 215 220
 Leu His Val Leu Ser Gly Met Lys Arg Lys Leu Leu Asn Ala His Leu
 225 230 235 240
 Lys Val Asp Ala Ile Val Glu Asp Val Ile Asn Glu His Lys Lys Asn
 245 250 255
 Leu Ala Thr Gly Lys Thr Asn Gly Ala Leu Gly Glu Asp Met Phe Ala
 260 265 270
 Ala Gly Thr Glu Thr Ser Ser Thr Thr Thr Val Trp Ala Met Ala Glu
 275 280 285
 Met Met Lys Asn Pro Asn Val Phe Asn Lys Ala Gln Ala Glu Val Arg
 290 295 300
 Glu Thr Phe Lys Asp Lys Val Thr Phe Asp Glu Ile Asp Ala Glu Glu
 305 310 315 320
 Leu Glu Tyr Leu Lys Leu Val Ile Lys Glu Thr Leu Arg Leu His Pro
 325 330 335
 Pro Ser Pro Leu Leu Val Pro Arg Glu Cys Arg Glu Asp Thr Asp Ile
 340 345 350
 Asn Gly Tyr Thr Ile Pro Ala Lys Thr Lys Val Met Val Asn Val Trp
 355 360 365
 Ala Leu Gly Arg Asp Pro Lys Tyr Trp Asp Asp Ala Glu Ser Phe Lys
 370 375 380
 Pro Glu Arg Phe Glu Gln Cys Ser Val Asp Phe Phe Gly Asn Asn Phe
 385 390 395 400
 Glu Phe Leu Pro Phe Gly Gly Gly Arg Arg Ile Cys Pro Gly Met Ser
 405 410 415
 Phe Gly Leu Ala Asn Leu Tyr Leu Pro Leu Ala Gln Leu Leu Tyr His
 420 425 430
 Phe Asp Trp Lys Leu Pro Ser Gly Met Met Pro Gly Asp Leu Asp Leu
 435 440 445
 Thr Glu Leu Ala Gly Ile Thr Ile Ala Arg Lys Gly Asp Leu Tyr Leu
 450 455 460
 Met Ala Thr Pro Tyr Gln Pro Ser Arg Glu
 465 470

<210> 4

<211> 1614

<212> DNA

<213> Nicotiana tabacum

<400> 4

ggatggtcta ataatcctcc atttatctcc caaaatggaa ttcttcagct tggtttccat 60
 attcctattc ctatctttcc tctttttggt aaggaaatgt aagaactcca atagccaaac 120
 caaacaattg cctccagggtc catggaaaat accaatacta ggaagtatgc ttcatatgct 180
 tgggtggagaa ccacaccata tccttaggga tttagccaaa aaatatggac caattatgca 240
 ccttcagttt ggtgaaattt ctgcagttgt ggttacttct agggagatgg caaaagaagt 300
 gctaaaaaact catgacgtag tttttgcata taggcctaaa attgtggcca tggacattat 360
 ctgttataac cagtctgata tcgccttttag cccttatggc gatcactgga gacaaatgcg 420

```

taaaatttgt gtcattggaac ttcttaatgc aaagaatggt cggctctttca gctcgatcag 480
acgtgatgaa gtcgttcgtc tcattgactc tattcgatca gattcttctt ctgggtgagct 540
agttaatttt acgcaaagga tcatttggtt cgcgagctcc atgacgtgta gatcagcatt 600
tggaagagta cttaaggggc aagacgtatt tgccaaaaag attagagaag taatagggtt 660
agcagaaggc tttgatgtgg ccgatatctt cccttcatac aagtttcttc atgttctcag 720
tggaatgaag cgtaaacttc tgaatgcca ccttaaggta gatgccattg ttgaggatgt 780
catcaacgag cacaagaaaa atcttgcaac tgggaaaact aatggagcat taggagacat 840
gtttgctgcc ggaacagaaa ctcatcaac aacaactgta tgggctatgg ctgaaatgat 900
gaagaatcca aatgtattca acaaagctca ggcagaagtg agagaaacct ttaaagacaa 960
agtaacattt gatgaaattg atgcagagga gctggaatac ttaaagttag ttattaaaga 1020
aactttgaga ctcatccac cgtctccact tttggtccca agagaatgta gggaagatac 1080
agatatcaac ggctacacta ttcctgcgaa gaccaaaagt atgggtaatg tttgggcatt 1140
gggaagagat ccaaaatatt gggatgacgc agaaaagctt aagccagaga gatttgagca 1200
atgctctgtg gatttttttg gtaataattt tgagtttctt ccctttggcg gtggacggag 1260
aatatgtcct ggtatgtcat ttggttttag taatctttac ttgccattgg ctcaattgct 1320
atatcacttt gattggaac tcccagcgg aatgatgccc ggagacttgg acttgactga 1380
attagctgga ataacaattg ctagaaaggg tgacctttac ttaatggcta ctccttatca 1440
accttctcgc gaataattta atggcatcag gttttttaat tccattgtca acctcactat 1500
tgtacaagct ttctgatgtt tcagggtttg ccgatttgta ataaatgtag tttttataat 1560
atgtatcata cccatgtaga agaggacga ttaattagtt gtaaaaaaaaa aaaa 1614

```

<210> 5

<211> 537

<212> PRT

<213> *Nicotiana tabacum*

<400> 5

```

Met Lys Asn Met Ala Lys Leu Leu Asn Lys Thr Ile Phe Cys Ile Leu
1          5          10          15
Phe Thr Ile Ala Phe Leu Ser Phe Ala Lys Leu Leu Ser Ser Tyr Leu
20          25          30
Ser Met Pro Phe Pro Leu Lys Tyr Met Ser Leu Ile Val Pro Leu Leu
35          40          45
Pro Leu Ile Ile Asn Phe Leu Tyr Val Lys Pro Gln Asn Asn Leu Pro
50          55          60
Pro Gly Pro Thr Ala Val Pro Ile Phe Gly Asn Trp Leu Gln Val Gly
65          70          75          80
Asn Asp Leu Asn His Gln Leu Leu Ala Thr Met Ser Gln Thr Tyr Gly
85          90          95
Pro Ile Phe Leu Leu Lys Leu Gly Ser Lys Asn Leu Ala Val Val Ser
100         105         110
Asn Pro Glu Leu Ala Asp Gln Val Leu His Thr Gln Gly Val Glu Phe
115         120         125
Gly Ser Arg Pro Arg Asn Val Val Phe Asp Ile Phe Thr Gly Asn Gly
130         135         140
Gln Asp Met Val Phe Thr Ile Tyr Gly Asp His Trp Arg Lys Met Arg
145         150         155         160
Arg Ile Met Thr Leu Pro Phe Phe Thr Asn Lys Val Val His Gln Tyr
165         170         175
Ser Asp Met Trp Glu Asn Glu Met Asp Leu Val Val Asn Asp Leu Lys
180         185         190

```



```

Lys Asn Glu Lys Val Lys Tyr Glu Gly Ile Val Ile Arg Lys Arg Leu
  195                200                205
Gln Leu Met Leu Tyr Asn Ile Met Tyr Arg Met Met Phe Asp Ala Lys
  210                215                220
Phe Glu Ser Gln Asn Asp Pro Leu Phe Ile Glu Ala Thr Lys Phe Asn
  225                230                235                240
Ser Glu Arg Ser Arg Leu Ala Gln Ser Phe Asp Tyr Asn Tyr Gly Asp
                245                250                255
Phe Ile Pro Leu Leu Arg Pro Phe Leu Arg Gly Tyr Leu Asn Lys Cys
                260                265                270
Lys Asp Leu Gln Thr Arg Arg Leu Ala Phe Phe Asn Asn Tyr Phe Val
  275                280                285
Glu Lys Arg Arg Lys Ile Met Asp Glu Asn Gly Glu Lys His Lys Ile
  290                295                300
Ser Cys Ala Ile Asp His Ile Ile Asp Ala Glu Met Lys Gly Glu Ile
  305                310                315                320
Asn Glu Gln Asn Val Leu Tyr Ile Val Glu Asn Ile Asn Val Ala Ala
                325                330                335
Ile Glu Thr Thr Leu Trp Ser Met Glu Trp Ala Ile Ala Glu Leu Val
  340                345                350
Asn His Pro Ile Val Gln Gln Lys Ile Arg Asp Glu Ile Ser Thr Val
  355                360                365
Leu Lys Gly Arg Ser Val Thr Glu Ser Asn Leu His Glu Leu Pro Tyr
  370                375                380
Leu Leu Ala Thr Val Asn Glu Thr Leu Arg Leu His Thr Pro Ile Pro
  385                390                395                400
Leu Leu Val Pro His Met Asn Leu Glu Glu Ala Lys Leu Gly Gly Tyr
                405                410                415
Thr Ile Pro Lys Glu Thr Lys Val Val Val Asn Ala Trp Trp Leu Ala
  420                425                430
Asn Asn Pro Ala Trp Trp Lys Asn Pro Asn Glu Phe Arg Pro Glu Arg
  435                440                445
Phe Leu Glu Glu Asp Ser Ser Thr Glu Ala Ala Val Ala Gly Gly Lys
  450                455                460
Val Asp Phe Arg Tyr Leu Pro Phe Gly Met Gly Arg Arg Ser Cys Pro
  465                470                475                480
Gly Ile Ile Leu Ala Leu Pro Ile Leu Gly Leu Val Ile Ala Lys Leu
                485                490                495
Val Ser Asn Phe Glu Met Gln Gly Pro Pro Gly Val Glu Lys Val Asp
  500                505                510
Thr Ser Glu Arg Gly Gly Gln Phe Ser Leu His Ile Ala Lys His Ser
  515                520                525
Thr Val Val Phe Lys Pro Ile Ala Ala
  530                535

```

<210> 6

<211> 1745

<212> DNA

<213> Nicotiana tabacum

```

<400> 6
cctctagcta atgaaaaaca tggccaaact tctcaacaag accatctttt gcattctctt 60
tacaattgca tttctttcat ttgccaaagt actgtcctcc tacctatcta tgcctttccc 120
tcttaagtac atgtcactta ttgtcccttt acttcccctt ataatacaact tcctctatgt 180
taagcccca aacaacctcc cacctgggtcc aacagcagtc ccaatatttg gtaattggct 240
tcaagttggc aatgacttga accatcaact ccttgccacc atgtcacaaa cctacgggtcc 300
tatattttta ctcaaacttg gttcaaaaaa cctagctgtg gtatcaaacc cagagctagc 360
tgaccaagtt ctacacacac aaggggtcga gtttgggtcc cgtccacgta acgttgtctt 420
cgacatatct actggtaatg gacaagacat ggtgttcacc atttatgggtg accattggcg 480
aaaaatgagg cgtattatga cgcttccatt tttcactaac aaagtgggtgc accaatatag 540
tgatatgtgg gagaatgaga tggacttagt tgtaatgac ttgaagaaga atgaaaaagt 600
gaaatatgag ggaattgtga ttaggaaacg attgcagctg atgctgtata acatcatgta 660
tcgaatgatg tttgatgcca aatttgagtc ccaaaatgat cctttgttca ttgaggcaac 720
aaagtttaat tcagagagaa gcagattagc tcagagcttt gactacaatt atggtgattt 780
tatcccttta cttagaccat tcttgagagg gtaccttaac aagtgtaaag acttacaaac 840
aaggagactt gcattcttca acaattattt tgtagagaaa agaaggaaaa taatggatga 900
aaatggagaa aagcataaga taagctgtgc tattgatcac attatagatg ccgaaatgaa 960
aggagaaata aatgagcaaa atgtactcta tattgtggag aatatcaatg ttgcagcaat 1020
tgaaacaact ctatggtcca tggaatgggc catagctgaa cttgtaaatc atcccattgt 1080
tcaacagaag attagggatg aaatctcaac agtcctcaa ggcagatcag tcacagaatc 1140
aaacctccat gagctgcctt acttgctagc aacagtaaat gaaacattaa gactccacac 1200
accaatacct ttacttgtac cccatatgaa ccttgaagaa gcaaagttag gtggttacac 1260
tattcctaaa gaaactaagg tggttgtgaa tgcgtgggtg ctggctaaca accctgcttg 1320
gtggaaaaac ccgaatgaat tccggcccga gaggtttctt gaggaggata gtagcacaga 1380
ggcagctgtt gctggtggca aggtagattt caggactta ccattcggta tggggaggcg 1440
gagctgcccc ggaatcatcc ttgcactgcc aattctgggg cttgtcatag ccaaactggt 1500
gtcaaatctt gaaatgcagg gtccaccagg tgtggaaaag gttgatacaa gtgaaagagg 1560
agggcagttt agcttgacac ttgcaaaaaca ttccacgggt gtcttcaagc ctattgtctg 1620
ataataatat gcttaagcta tccttgtttt aattatattt gtcttaccag aaagcaaaac 1680
tactaagtta ctcgataaag atttcaatga atattacagt ttttgttaaa aaaaaaaaaa 1740
aaaaa

```

<210> 7

<211> 534

<212> PRT

<213> Nicotiana tabacum

<400> 7

```

Met Ala Lys Leu Leu Asn Asn Thr Ile Phe Cys Ile Leu Phe Ser Ile
  1             5             10             15
Val Phe Leu Ser Phe Ala Lys Leu Leu Ser Ser Tyr Leu Ser Ile Pro
      20             25             30
Phe Pro Leu Glu Tyr Ile Ser Leu Ile Val Leu Leu Leu Pro Leu Ile
      35             40             45
Ile Asn Phe Leu Cys Val Lys Pro Gln Asn Asn Leu Pro Pro Gly Pro
      50             55             60
Thr Ala Val Pro Ile Phe Gly Asn Trp Leu Gln Val Gly Asn Asp Leu
      65             70             75             80
Asn His Gln Leu Leu Ala Thr Met Ser Gln Thr Tyr Gly Pro Ile Phe
      85             90             95
Leu Leu Lys Leu Gly Ser Lys Asn Leu Ala Val Val Ser Asn Pro Glu

```

			100						105						110		
Leu	Ala	Asn	Gln	Val	Leu	His	Thr	Gln	Gly	Val	Glu	Phe	Gly	Ser	Arg		
		115						120				125					
Pro	Arg	Asn	Val	Val	Phe	Asp	Ile	Phe	Thr	Gly	Asn	Gly	Gln	Asp	Met		
		130					135				140						
Val	Phe	Thr	Ile	Tyr	Gly	Asp	His	Trp	Arg	Lys	Met	Arg	Arg	Ile	Met		
145					150					155					160		
Thr	Leu	Pro	Phe	Phe	Thr	Asn	Lys	Val	Val	His	Gln	Tyr	Ser	Asp	Met		
				165					170					175			
Trp	Glu	Asn	Glu	Met	Asp	Leu	Val	Val	Asp	Asp	Leu	Lys	Lys	Asn	Glu		
			180					185					190				
Lys	Val	Lys	Tyr	Asp	Gly	Ile	Val	Ile	Arg	Lys	Arg	Leu	Gln	Leu	Met		
		195					200					205					
Leu	Tyr	Asn	Ile	Met	Tyr	Arg	Met	Met	Phe	Asp	Ala	Lys	Phe	Glu	Ser		
		210					215				220						
Gln	Asp	Asp	Pro	Leu	Phe	Ile	Glu	Ala	Thr	Lys	Phe	Asn	Ser	Glu	Arg		
225					230					235					240		
Ser	Arg	Leu	Ala	Gln	Ser	Phe	Asp	Tyr	Asn	Tyr	Gly	Asp	Phe	Ile	Pro		
				245					250					255			
Leu	Leu	Arg	Pro	Phe	Leu	Lys	Gly	Tyr	Leu	Asn	Lys	Cys	Lys	Asp	Leu		
			260					265					270				
Gln	Thr	Arg	Arg	Leu	Ala	Phe	Phe	Asn	Asn	Tyr	Phe	Val	Gly	Lys	Arg		
		275						280				285					
Arg	Lys	Ile	Met	Gly	Glu	Asn	Gly	Glu	Lys	His	Lys	Ile	Cys	Cys	Ala		
		290					295				300						
Ile	Asp	His	Ile	Ile	Asp	Ala	Glu	Met	Lys	Gly	Glu	Ile	Ser	Glu	Gln		
305					310					315					320		
Asn	Val	Leu	Tyr	Ile	Val	Glu	Asn	Ile	Asn	Val	Ala	Ala	Ile	Glu	Thr		
				325					330					335			
Thr	Leu	Trp	Ser	Met	Glu	Trp	Ala	Ile	Ala	Glu	Leu	Val	Asn	His	Pro		
			340					345					350				
Ile	Val	Gln	Gln	Lys	Ile	Arg	Asp	Glu	Ile	Ser	Thr	Val	Leu	Lys	Gly		
		355					360					365					
Lys	Ser	Val	Lys	Glu	Ser	Asn	Leu	His	Glu	Leu	Pro	Tyr	Leu	Leu	Ala		
		370					375				380						
Thr	Val	Asn	Glu	Thr	Leu	Arg	Leu	His	Thr	Pro	Ile	Pro	Leu	Leu	Val		
385					390					395					400		
Pro	His	Met	Asn	Leu	Glu	Glu	Ala	Lys	Leu	Gly	Gly	Tyr	Thr	Ile	Pro		
				405					410					415			
Lys	Glu	Thr	Lys	Val	Val	Val	Asn	Ala	Trp	Trp	Leu	Ala	Asn	Asn	Pro		
			420					425					430				
Ala	Trp	Trp	Lys	Asn	Gln	Asn	Glu	Phe	Arg	Pro	Glu	Arg	Phe	Leu	Glu		
		435					440										

```
<210> 8
<211> 1693
<212> DNA
<213> Nicotiana tabacum
```

```
<210> 9
<211> 519
<212> PRT
<213> Nicotiana tabacum
```

- 9 -

- 10 -

```

Thr Thr Asn Ala Ala Ile Asp Tyr Arg Gly Gln His Tyr Glu Leu Ile
      435      440      445
Pro Phe Gly Ser Gly Arg Arg Ala Cys Pro Ala Met Asn Tyr Ser Leu
      450      455      460
Gln Val Glu His Leu Ser Ile Ala His Leu Ile Gln Gly Phe Asn Phe
465      470      475      480
Ala Thr Thr Thr Asn Glu Pro Leu Asp Met Lys Gln Gly Val Gly Leu
      485      490      495
Thr Leu Pro Lys Lys Thr Asp Val Glu Val Leu Ile Thr Pro Arg Leu
      500      505      510
Pro Pro Thr Leu Tyr Gln Tyr
      515

```

<210> 10

<211> 1578

<212> DNA

<213> Nicotiana tabacum

<400> 10

```

atgtatcatc ttctttctcc catagaagcc attgtaggac ttgtaacctt tgcatttcta 60
ctctacttgc tatggacaaa aaaacaatca aaaatcttaa acccactgcc tccaaaaatc 120
ccaggtggat ggccagtaat cggccatctc ttttatttca acaacaatgg cgatgatgac 180
cgccattttt ctcaaaaact cggagactta gctgacaaat atgggtcccg cttcacattc 240
cggttagggt ttgcgcggtt cttggcgggt agtagttatg aagctatgaa agaatgcttc 300
tctaccaatg atatccattt cgccgatcgg ccagctttac tttacggaga atacctttgc 360
tataacaatg ccatgcttgc tgttgccaaa tatggccctt actggaaaaa aaatcgaaag 420
ctagtcaatc aagaacttct ctccgttagt cggctcgaaa aattcaaaca tgtagattt 480
tctatagttc agaaaaatat taaacaattg tataattgtg attcaccaat ggtgaagata 540
aaccttagtg attggataga taaattgaca ttcgacatca ttttgaaaat ggttggtggg 600
aagacctata ataattggaca tggagaaata ctcaaagcag cttttcagaa gttcatggtt 660
caagctatgg agattgagct ctatgatgtt tttcacattc catttttcaa gtggttggat 720
cttacaggga atattaaggc tatgaaacaa actttcaaag acattgataa tattatccaa 780
ggttggttag atgagcacat taagaagaga gaaacaaagg atgttgaggg tgaaaatgaa 840
caagatttta ttgatgtgct gctttccaag aggagcaacg aacatcttgg cgatgggttac 900
tctcatgaca ccaccatcaa agcaacagta ttcactttgg tcttggtatg aacagacaca 960
cttgcacttc atataaagtg ggtaatggcg ttaatgataa acaataagaa tgatcatgaag 1020
aaagcacaa gagagatgga caccattgtt ggtagagata gatgggtaga agagaatgat 1080
atcaagaatt tgggtgtatc tcaagcaatt gttaaagaag tattacgatt acatccacct 1140
gcacctttgt cagtacaaca cctatccgta aaagattgtg ttgtcaatgg ataccatatt 1200
cctaagggga ctgcactact tacaaaatatt atgaaacttc aacgagaccc acaaatatgg 1260
gtagatcctg atacattcga tccagaaaga ttcttgacga ctaatgctgc aattgactat 1320
cgcgggcagc actatgagtt gatcccgtt ggatcaggga gacgagcttg tcccgcgatg 1380
aattactcat tgcaagtgga acacctttca attgctcatt tgatccaggg tttcaatttt 1440
gcaactacga ctaacgagcc tttggatatg aaacaaggcg tgggtctaac tttacctaa 1500
aagacagatg ttgaagtgtt aattacacct cgccttcctc ctacgcttta tcaatattaa 1560
tatgttttgt tgttgtga
1578

```

<210> 11

<211> 509

<212> PRT

<213> Nicotiana tabacum

<400> 11

```

Met Glu Gly Thr Asn Leu Thr Thr Tyr Ala Ala Val Phe Leu Gly Thr
 1           5           10           15
Leu Phe Leu Leu Leu Leu Ser Lys Phe Leu Arg Gln Arg Lys Leu Asn
          20           25           30
Leu Pro Pro Gly Pro Lys Pro Trp Pro Ile Ile Gly Asn Leu Asn Leu
          35           40           45
Ile Gly Asn Leu Pro His Arg Ser Ile His Glu Leu Ser Leu Lys Tyr
 50           55           60
Gly Pro Ile Met Gln Leu Gln Phe Gly Thr Phe Pro Val Val Val Gly
65           70           75           80
Ser Ser Val Glu Met Ala Lys Val Phe Leu Lys Ser Met Asp Ile Asn
          85           90           95
Phe Val Gly Arg Pro Lys Thr Ala Ala Gly Lys Tyr Thr Thr Tyr Asn
          100          105          110
Tyr Ser Asp Ile Thr Trp Ser Pro Tyr Gly Pro Tyr Trp Arg Gln Ala
          115          120          125
Arg Arg Met Cys Leu Met Glu Leu Phe Ser Thr Lys Arg Leu Asp Ser
          130          135          140
Tyr Glu Tyr Ile Arg Ala Glu Glu Leu His Ser Leu Leu His Asn Leu
          145          150          155          160
Asn Lys Ile Ser Gly Lys Pro Ile Val Leu Lys Asp Tyr Leu Thr Thr
          165          170          175
Leu Ser Leu Asn Val Ile Ser Arg Met Val Leu Gly Lys Arg Tyr Leu
          180          185          190
Asp Glu Ser Glu Asn Ser Ile Val Thr Pro Glu Glu Phe Lys Lys Met
          195          200          205
Leu Asp Glu Leu Phe Leu Leu Asn Gly Val Leu Asn Ile Gly Asp Ser
          210          215          220
Ile Pro Trp Ile Asp Phe Met Asp Leu Gln Gly Tyr Val Lys Arg Met
          225          230          235          240
Lys Phe Val Ser Lys Lys Phe Asp Lys Phe Leu Glu His Val Ile Asp
          245          250          255
Glu His Asn Val Arg Arg Asn Gly Val Glu Asn Tyr Ile Ala Lys Asp
          260          265          270
Met Val Asp Val Leu Leu Gln Leu Ala Asp Asp Pro Thr Leu Glu Val
          275          280          285
Lys Leu Glu Arg His Gly Val Lys Ala Phe Thr Gln Asp Met Leu Ala
          290          295          300
Gly Gly Thr Glu Ser Ser Ala Val Thr Val Glu Trp Ala Ile Ser Glu
          305          310          315          320
Leu Leu Lys Lys Pro Glu Ile Phe Lys Lys Ala Thr Glu Glu Leu Asp
          325          330          335
Arg Val Ile Gly Gln Asn Arg Trp Val Gln Glu Lys Asp Ile Pro Asn
          340          345          350
Leu Pro Tyr Ile Glu Ala Ile Val Lys Glu Thr Met Arg Leu His Pro
          355          360          365
Val Ala Pro Met Leu Val Pro Arg Glu Cys Arg Glu Asp Cys Lys Val
          370          375          380

```

Ala	Gly	Tyr	Asp	Val	Lys	Lys	Gly	Thr	Arg	Val	Leu	Val	Ser	Val	Trp
385					390					395					400
Thr	Ile	Gly	Arg	Asp	Pro	Thr	Leu	Trp	Asp	Glu	Pro	Glu	Ala	Phe	Lys
			405						410					415	
Pro	Glu	Arg	Phe	His	Glu	Lys	Ser	Ile	Asp	Val	Lys	Gly	His	Asp	Phe
			420					425					430		
Glu	Leu	Leu	Pro	Phe	Gly	Ala	Gly	Arg	Arg	Met	Cys	Pro	Gly	Tyr	Asn
	435						440					445			
Leu	Gly	Leu	Lys	Val	Ile	Gln	Ala	Ser	Leu	Ala	Asn	Leu	Ile	His	Gly
	450					455					460				
Phe	Asn	Trp	Ser	Leu	Pro	Asp	Asn	Met	Thr	Pro	Glu	Asp	Leu	Asp	Met
465					470					475					480
Asp	Glu	Ile	Phe	Gly	Leu	Ser	Thr	Pro	Lys	Lys	Phe	Pro	Leu	Ala	Thr
			485						490						495
Val	Ile	Glu	Pro	Arg	Leu	Ser	Pro	Lys	Leu	Tyr	Ser	Val			
			500					505							

<210> 12

<211> 1530

<212> DNA

<213> Nicotiana tabacum

<400> 12

```

atggaaggta caaacttgac tacatatgca gcagtatttc ttggtactct gtttcttttg 60
ctcctttcca aatttcttcg ccaaagaaaa ctcaacttac ctccaggccc aaaaccatgg 120
ccgatcatcg gaaacttaaa ccttatcggc aatcttcctc atcgctcaat ccacgaactt 180
tactcaagt acgggccaat tatgcaactc caattcggga ctttccccgt tgttgttggc 240
tcttcgctcg aaatggccaa ggttttcctc aaatcaatgg atattaactt tgtaggcagg 300
cctaaaacgg ccgcccggaa gtacacaact tacaattatt cagatattac atggtctcct 360
tatggaccat attggcgcca ggcacgtaga atgtgcctaa tgggaattatt cagcacgaaa 420
cgtctcgatt catacgagta tattcgggct gaggagttgc attctctgct ccataatttg 480
aataaaatat cagggaaacc aattgtgctg aaagattatt tgacgacggt gagtttaaat 540
gttattagca ggatggtact ggggaagagg tatttggacg aatccgagaa ctcgattgtg 600
actcctgagg aatttaagaa gatgttggac gagctgttct tgctaaatgg tgtacttaat 660
attggagatt caattccctg gattgatttc atggacttac aaggttatgt taagaggatg 720
aaatttgtga gcaagaaatt cgacaagttt ttggagcatg ttatcgatga gcataacggt 780
aggagaaatg gagtggagaa ttacattgct aaggacatgg ttgatgttct gttgcagctt 840
gctgatgac cgacgttgga agttaagctg gagagacatg gagtcaaagc attcactcag 900
gatatgcttg ctggtggaac cgagagttca gcagtgcacg tggagtgggc aatttcggag 960
ctgctaaaga agccagagat tttcaaaaag gctacagaag aattggatcg agtaattggg 1020
cagaatagat ggttacaaga aaaagacatt ccaaactctc cttacataga ggcaatagtc 1080
aaagagacta tgcgactgca ccccgtagga ccaatgttgg tgccacggga gtgtcgagaa 1140
gactgtaagg tagcaggcta cgacgttaag aaaggaacca gggtccttgt gagcgtatgg 1200
actattggaa gagaccctac attgtgggac gagcctgagg cgttcaagcc ggagaggttc 1260
cacgaaaagt ccattgatgt taaaggacat gattttgagc ttttgccatt tggagctggg 1320
agaaggatgt gcccggtta taacttgggg cttaagggtga ttcaagctag cttagctaata 1380
cttatacatg gatttaactg gtcattgcct gataaatatga ctcctgagga cctcgacatg 1440
gatgagattt ttgggctctc cacacctaaa aagtttccac ttgctactgt gattgagcca 1500
agactttcac caaaacttta ctctgtttga 1530

```


<210> 13
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 13
ggcggagaat ttgtcctgga atgtcatttg gtttag 36

<210> 14
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 14
gtacaatagt gaggttgaca atg 23

<210> 15
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 15
ggtggttgatg aatgcatg 18

<210> 16
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 16
ttatgcagca ataggcttga agaca 25

<210> 17
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 17
gggggatcca tgcaattctt cagcttggtt tcc 33

<210> 18
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 18
ggggaattct tactctcgag aaggttgata agg 33

<210> 19
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 19
cccggatcca tgtatcatct tctttctccc 30

<210> 20
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 20
ggggaattct caatattgat aaagcgtagg agg 33

<210> 21
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 21
cccggatcca tgcaatcctt cagcttggtt tcc 33

<210> 22
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 22
ggggagctct cactcgcaag aagattgata agg 33

<210> 23
<211> 75
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 23
gccattatcg gcgcaataact aatctccaaa ctccgcggta aaaaattcaa gctcccacct 60
ggtccaacag cagtc 75

<210> 24
<211> 75
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 24
gggggatcca tggacctcct cctcatagaa aaaaccctcg tcgccttatt cgccgccatt 60
atcggcgcaa tacta 75

<210> 25
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 gene

<400> 25
ggggagctct tatgcagcaa taggcttgaa gac 33

<210> 26
<211> 7
<212> PRT
<213> Nicotiana tabacum p450 protein

<220>
<221> VARIANT
<222> 7
<223> Xaa = Any Amino Acid

<400> 26

Lys Glu Thr Leu Arg Leu Xaa
1 5

<210> 27

<211> 11

<212> PRT

<213> Nicotiana tabacum p450 protein

<220>

<221> VARIANT

<222> 4,8

<223> Xaa = Any Amino Acid

<400> 27

Pro Phe Gly Xaa Gly Arg Arg Xaa Cys Pro Ala
1 5 10

<210> 28

<211> 11

<212> PRT

<213> Nicotiana tabacum p450 protein

<220>

<221> VARIANT

<222> 4,8

<223> Xaa = Any Amino Acid

<400> 28

Pro Phe Gly Xaa Gly Arg Arg Xaa Cys Pro Gly
1 5 10

<210> 29

<211> 6

<212> PRT

<213> Nicotiana tabacum p450 protein

<220>

<221> VARIANT

<222> 2

<223> Xaa = Any Amino Acid

<400> 29

Phe Xaa Pro Glu Arg Phe
1 5

<210> 30

<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 protein

<400> 30
Ala Ala Arg Gly Ala Arg Ala Cys Ile Tyr Thr Ile Met Gly Ile Tyr
1 5 10 15
Thr Ile Cys Ala
20

<210> 31
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 protein

<400> 31
Ala Ala Arg Gly Ala Arg Ala Cys Ile Tyr Thr Ile Met Gly Ile Tyr
1 5 10 15
Thr Ile Thr Ala
20

<210> 32
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 protein

<400> 32
Ala Ala Arg Gly Ala Arg Ala Cys Ile Tyr Thr Ile Met Gly Ile Tyr
1 5 10 15
Thr Ile Met Gly
20

<210> 33
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> derived from Nicotiana tabacum p450 protein

<400> 33

Thr Thr Tyr Ile Ile Ile Cys Cys Ile Gly Ala Arg Met Gly Ile Thr
1 5 10 15
Thr Tyr

<210> 34

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> derived from Nicotiana tabacum p450 protein

<400> 34

Arg Ala Ala Ile Cys Lys Tyr Thr Cys Ile Gly Gly Ile Ile Ile Arg
1 5 10 15
Ala Ala

<210> 35

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> derived from Nicotiana tabacum p450 protein

<400> 35

Gly Gly Ile Met Gly Ile Met Gly Ile Ile Ile Ile Thr Gly Tyr Cys
1 5 10 15
Cys Ile Gly Ser
20

<210> 36

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> derived from Nicotiana tabacum p450 protein

<400> 36

Cys Lys Ile Cys Lys Ile Cys Cys Ile Ile Ile Ile Cys Cys Arg Ala
1 5 10 15
Ala Ile Gly Gly
20

<210> 37
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> derived from T7 Bacteriophage Promoter

<400> 37
 gtaatacgac tcactatagg g

21

<210> 38
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> derived from T3 Bacteriophage Promoter

<400> 38
 caattaaccc tcactaaagg g

21

<210> 39
 <211> 500
 <212> PRT
 <213> Mentha piperita

<400> 39
 Met Glu Leu Gln Ile Ser Ser Ala Ile Ile Ile Leu Val Val Thr Tyr
 1 5 10 15
 Thr Ile Ser Leu Leu Ile Ile Lys Gln Trp Arg Lys Pro Lys Pro Gln
 20 25 30
 Glu Asn Leu Pro Pro Gly Pro Pro Lys Leu Pro Leu Ile Gly His Leu
 35 40 45
 His Leu Leu Trp Gly Lys Leu Pro Gln His Ala Leu Ala Ser Val Ala
 50 55 60
 Lys Gln Tyr Gly Pro Val Ala His Val Gln Leu Gly Glu Val Phe Ser
 65 70 75 80
 Val Val Leu Ser Ser Arg Glu Ala Thr Lys Glu Ala Met Lys Leu Val
 85 90 95
 Asp Pro Ala Cys Ala Asp Arg Phe Glu Ser Ile Gly Thr Lys Ile Met
 100 105 110
 Trp Tyr Asp Asn Asp Asp Ile Ile Phe Ser Pro Tyr Ser Val His Trp
 115 120 125
 Arg Gln Met Arg Lys Ile Cys Val Ser Glu Leu Leu Ser Ala Arg Asn
 130 135 140
 Val Arg Ser Phe Gly Phe Ile Arg Gln Asp Glu Val Ser Arg Leu Leu
 145 150 155 160
 Gly His Leu Arg Ser Ser Ala Ala Ala Gly Glu Ala Val Asp Leu Thr
 165 170 175
 Glu Arg Ile Ala Thr Leu Thr Cys Ser Ile Ile Cys Arg Ala Ala Phe

```
<210> 40
<211> 496
<212> PRT
<213> Mentha spicata
```

-21-


```

Ile Val Ser Leu Leu Ile Asn Gln Trp Arg Lys Ser Lys Ser Gln Gln
      20      25      30
Asn Leu Pro Pro Ser Pro Pro Lys Leu Pro Val Ile Gly His Leu His
      35      40      45
Phe Leu Trp Gly Gly Leu Pro Gln His Val Phe Arg Ser Ile Ala Gln
      50      55      60
Lys Tyr Gly Pro Val Ala His Val Gln Leu Gly Glu Val Tyr Ser Val
      65      70      75      80
Val Leu Ser Ser Ala Glu Ala Ala Lys Gln Ala Met Lys Val Leu Asp
      85      90      95
Pro Asn Phe Ala Asp Arg Phe Asp Gly Ile Gly Ser Arg Thr Met Trp
      100      105      110
Tyr Asp Lys Asp Asp Ile Ile Phe Ser Pro Tyr Asn Asp His Trp Arg
      115      120      125
Gln Met Arg Arg Ile Cys Val Thr Glu Leu Leu Ser Pro Lys Asn Val
      130      135      140
Arg Ser Phe Gly Tyr Ile Arg Gln Glu Glu Ile Glu Arg Leu Ile Arg
      145      150      155      160
Leu Leu Gly Ser Ser Gly Gly Ala Pro Val Asp Val Thr Glu Glu Val
      165      170      175
Ser Lys Met Ser Cys Val Val Val Cys Arg Ala Ala Phe Gly Ser Val
      180      185      190
Leu Lys Asp Gln Gly Ser Leu Ala Glu Leu Val Lys Glu Ser Leu Ala
      195      200      205
Leu Ala Ser Gly Phe Glu Leu Ala Asp Leu Tyr Pro Ser Ser Trp Leu
      210      215      220
Leu Asn Leu Leu Ser Leu Asn Lys Tyr Arg Leu Gln Arg Met Arg Arg
      225      230      235      240
Arg Leu Asp His Ile Leu Asp Gly Phe Leu Glu Glu His Arg Glu Lys
      245      250      255
Lys Ser Gly Glu Phe Gly Gly Glu Asp Ile Val Asp Val Leu Phe Arg
      260      265      270
Met Gln Lys Gly Ser Asp Ile Lys Ile Pro Ile Thr Ser Asn Cys Ile
      275      280      285
Lys Gly Phe Ile Phe Asp Thr Phe Ser Ala Gly Ala Glu Thr Ser Ser
      290      295      300
Thr Thr Ile Ser Trp Ala Leu Ser Glu Leu Met Arg Asn Pro Ala Lys
      305      310      315      320
Met Ala Lys Val Gln Ala Glu Val Arg Glu Ala Leu Lys Gly Lys Thr
      325      330      335
Val Val Asp Leu Ser Glu Val Gln Glu Leu Lys Tyr Leu Arg Ser Val
      340      345      350
Leu Lys Glu Thr Leu Arg Leu His Pro Pro Phe Pro Leu Ile Pro Arg
      355      360      365
Gln Ser Arg Glu Glu Cys Glu Val Asn Gly Tyr Thr Ile Pro Ala Lys
      370      375      380
Thr Arg Ile Phe Ile Asn Val Trp Ala Ile Gly Arg Asp Pro Gln Tyr
      385      390      395      400
Trp Glu Asp Pro Asp Thr Phe Arg Pro Glu Arg Phe Asp Glu Val Ser
      405      410      415
Arg Asp Phe Met Gly Asn Asp Phe Glu Phe Ile Pro Phe Gly Ala Gly

```

			420					425				430				
Arg	Arg	Ile	Cys	Pro	Gly	Leu	His	Phe	Gly	Leu	Ala	Asn	Val	Glu	Ile	
		435					440					445				
Pro	Leu	Ala	Gln	Leu	Leu	Tyr	His	Phe	Asp	Trp	Lys	Leu	Pro	Gln	Gly	
	450					455					460					
Met	Thr	Asp	Ala	Asp	Leu	Asp	Met	Thr	Glu	Thr	Pro	Gly	Leu	Ser	Gly	
465					470				475						480	
Pro	Lys	Lys	Lys	Asn	Val	Cys	Leu	Val	Pro	Thr	Leu	Tyr	Lys	Ser	Pro	
			485					490						495		

```
<210> 41
<211> 509
<212> PRT
<213> Nepeta racemosa
```

<400> 41																
Met	Val	Ser	Leu	Ser	Tyr	Phe	Leu	Ile	Ala	Leu	Leu	Cys	Thr	Leu	Pro	
1				5					10					15		
Phe	Leu	Leu	Phe	Leu	Asn	Lys	Trp	Arg	Arg	Ser	Tyr	Ser	Gly	Lys	Thr	
			20				25						30			
Pro	Pro	Pro	Ser	Pro	Pro	Lys	Leu	Pro	Val	Ile	Gly	Asn	Leu	His	Gln	
		35				40					45					
Leu	Gly	Leu	Tyr	Pro	His	Arg	Tyr	Leu	Gln	Ser	Leu	Ser	Arg	Arg	Tyr	
	50				55					60						
Gly	Pro	Leu	Met	Gln	Leu	His	Phe	Gly	Ser	Val	Pro	Val	Leu	Val	Ala	
65				70					75						80	
Ser	Ser	Pro	Glu	Ala	Ala	Arg	Glu	Ile	Met	Lys	Asn	Gln	Asp	Ile	Val	
			85				90						95			
Phe	Ser	Asn	Arg	Pro	Lys	Met	Ser	Ile	Ala	Asn	Arg	Leu	Phe	Phe	Asn	
		100				105						110				
Asn	Arg	Asp	Val	Ala	Phe	Thr	Gln	Tyr	Gly	Glu	Tyr	Trp	Arg	Gln	Ile	
		115				120					125					
Arg	Ser	Ile	Cys	Val	Leu	Gln	Leu	Leu	Ser	Asn	Lys	Arg	Val	Gln	Ser	
	130				135					140						
Phe	Arg	Arg	Val	Arg	Glu	Glu	Glu	Thr	Ser	Ile	Met	Val	Glu	Lys	Ile	
145				150					155						160	
Met	Gln	Leu	Gly	Ser	Ser	Ser	Ser	Thr	Pro	Val	Asn	Leu	Ser	Glu	Leu	
			165				170						175			
Leu	Leu	Ser	Leu	Thr	Asn	Asp	Val	Val	Cys	Arg	Val	Thr	Leu	Gly	Lys	
		180				185						190				
Lys	Tyr	Gly	Gly	Gly	Asn	Gly	Ser	Glu	Glu	Val	Asp	Lys	Leu	Lys	Glu	
		195				200				205						
Met	Leu	Thr	Glu	Ile	Gln	Asn	Leu	Met	Gly	Ile	Ser	Pro	Val	Trp	Glu	
	210				215					220						
Phe	Ile	Pro	Trp	Leu	Asn	Trp	Thr	Arg	Arg	Phe	Asp	Gly	Val	Asp	Gln	
225				230					235						240	
Arg	Val	Asp	Arg	Ile	Val	Lys	Ala	Phe	Asp	Gly	Phe	Leu	Glu	Ser	Val	
		245				250						255				
Ile	Gln	Glu	His	Lys	Glu	Arg	Asp	Gly	Asp	Lys	Asp	Gly	Asp	Gly	Asp	
		260				265						270				

Gly Ala Leu Asp Phe Val Asp Ile Leu Leu Gln Phe Gln Arg Glu Asn
 275 280 285
 Lys Asn Arg Ser Pro Val Glu Asp Asp Thr Val Lys Ala Leu Ile Leu
 290 295 300
 Asp Met Phe Val Ala Gly Thr Asp Thr Thr Ala Thr Ala Leu Glu Trp
 305 310 315 320
 Ala Val Ala Glu Leu Ile Lys Asn Pro Arg Ala Met Lys Arg Leu Gln
 325 330 335
 Asn Glu Val Arg Glu Val Ala Gly Ser Lys Ala Glu Ile Glu Glu Glu
 340 345 350
 Asp Leu Glu Lys Met Pro Tyr Leu Lys Ala Ser Ile Lys Glu Ser Leu
 355 360 365
 Arg Leu His Val Pro Val Val Leu Leu Val Pro Arg Glu Ser Thr Arg
 370 375 380
 Asp Thr Asn Val Leu Gly Tyr Asp Ile Ala Ser Gly Thr Arg Val Leu
 385 390 395 400
 Ile Asn Ala Trp Ala Ile Ala Arg Asp Pro Ser Val Trp Glu Asn Pro
 405 410 415
 Glu Glu Phe Leu Pro Glu Arg Phe Leu Asp Ser Ser Ile Asp Tyr Lys
 420 425 430
 Gly Leu His Phe Glu Leu Leu Pro Phe Gly Ala Gly Arg Arg Gly Cys
 435 440 445
 Pro Gly Ala Thr Phe Ala Val Ala Ile Asp Glu Leu Ala Leu Ala Lys
 450 455 460
 Leu Val His Lys Phe Asp Phe Gly Leu Pro Asn Gly Ala Arg Met Glu
 465 470 475 480
 Glu Leu Asp Met Ser Glu Thr Ser Gly Met Thr Val His Lys Lys Ser
 485 490 495
 Pro Leu Leu Leu Leu Pro Ile Pro His His Ala Ala Pro
 500 505

<210> 42

<211> 471

<212> PRT

<213> Persea americana

<400> 42

Met Ala Ile Leu Val Ser Leu Leu Phe Leu Ala Ile Ala Leu Thr Phe
 1 5 10 15
 Phe Leu Leu Lys Leu Asn Glu Lys Arg Glu Lys Lys Pro Asn Leu Pro
 20 25 30
 Pro Ser Pro Pro Asn Leu Pro Ile Ile Gly Asn Leu His Gln Leu Gly
 35 40 45
 Asn Leu Pro His Arg Ser Leu Arg Ser Leu Ala Asn Glu Leu Gly Pro
 50 55 60
 Leu Ile Leu Leu His Leu Gly His Ile Pro Thr Leu Ile Val Ser Thr
 65 70 75 80
 Ala Glu Ile Ala Glu Glu Ile Leu Lys Thr His Asp Leu Ile Phe Ala
 85 90 95
 Ser Arg Pro Ser Thr Thr Ala Ala Arg Arg Ile Phe Tyr Asp Cys Thr

<210> 43
<211> 509

<212> PRT

<213> Arabidopsis thaliana

<400> 43

```

Met Ala Phe Phe Ser Met Ile Ser Ile Leu Leu Gly Phe Val Ile Ser
 1           5           10           15
Ser Phe Ile Phe Ile Phe Phe Phe Lys Lys Leu Leu Ser Phe Ser Arg
           20           25           30
Lys Asn Met Ser Glu Val Ser Thr Leu Pro Ser Val Pro Val Val Pro
           35           40           45
Gly Phe Pro Val Ile Gly Asn Leu Leu Gln Leu Lys Glu Lys Lys Pro
 50           55           60
His Lys Thr Phe Thr Arg Trp Ser Glu Ile Tyr Gly Pro Ile Tyr Ser
 65           70           75           80
Ile Lys Met Gly Ser Ser Ser Leu Ile Val Leu Asn Ser Thr Glu Thr
           85           90           95
Ala Lys Glu Ala Met Val Thr Arg Phe Ser Ser Ile Ser Thr Arg Lys
           100          105          110
Leu Ser Asn Ala Leu Thr Val Leu Thr Cys Asp Lys Ser Met Val Ala
           115          120          125
Thr Ser Asp Tyr Asp Asp Phe His Lys Leu Val Lys Arg Cys Leu Leu
 130          135          140
Asn Gly Leu Leu Gly Ala Asn Ala Gln Lys Arg Lys Arg His Tyr Arg
 145          150          155          160
Asp Ala Leu Ile Glu Asn Val Ser Ser Lys Leu His Ala His Ala Arg
           165          170          175
Asp His Pro Gln Glu Pro Val Asn Phe Arg Ala Ile Phe Glu His Glu
           180          185          190
Leu Phe Gly Val Ala Leu Lys Gln Ala Phe Gly Lys Asp Val Glu Ser
           195          200          205
Ile Tyr Val Lys Glu Leu Gly Val Thr Leu Ser Lys Asp Glu Ile Phe
 210          215          220
Lys Val Leu Val His Asp Met Met Glu Gly Ala Ile Asp Val Asp Trp
 225          230          235          240
Arg Asp Phe Phe Pro Tyr Leu Lys Trp Ile Pro Asn Lys Ser Phe Glu
           245          250          255
Ala Arg Ile Gln Gln Lys His Lys Arg Arg Leu Ala Val Met Asn Ala
           260          265          270
Leu Ile Gln Asp Arg Leu Lys Gln Asn Gly Ser Glu Ser Asp Asp Asp
 275          280          285
Cys Tyr Leu Asn Phe Leu Met Ser Glu Ala Lys Thr Leu Thr Lys Glu
 290          295          300
Gln Ile Ala Ile Leu Val Trp Glu Thr Ile Ile Glu Thr Ala Asp Thr
 305          310          315          320
Thr Leu Val Thr Thr Glu Trp Ala Ile Tyr Glu Leu Ala Lys His Pro
           325          330          335
Ser Val Gln Asp Arg Leu Cys Lys Glu Ile Gln Asn Val Cys Gly Gly
           340          345          350
Glu Lys Phe Lys Glu Glu Gln Leu Ser Gln Val Pro Tyr Leu Asn Gly
           355          360          365
Val Phe His Glu Thr Leu Arg Lys Tyr Ser Pro Ala Pro Leu Val Pro

```

370		375		380
Ile Arg Tyr Ala His Glu Asp Thr Gln Ile Gly Gly Tyr His Val Pro				
385		390		395
Ala Gly Ser Glu Ile Ala Ile Asn Ile Tyr Gly Cys Asn Met Asp Lys				400
	405		410	415
Lys Arg Trp Glu Arg Pro Glu Asp Trp Trp Pro Glu Arg Phe Leu Asp				
	420		425	430
Asp Gly Lys Tyr Glu Thr Ser Asp Leu His Lys Thr Met Ala Phe Gly				
	435		440	445
Ala Gly Lys Arg Val Cys Ala Gly Ala Leu Gln Ala Ser Leu Met Ala				
	450		455	460
Gly Ile Ala Ile Gly Arg Leu Val Gln Glu Phe Glu Trp Lys Leu Arg				
465		470		475
Asp Gly Glu Glu Glu Asn Val Asp Thr Tyr Gly Leu Thr Ser Gln Lys				480
	485		490	495
Leu Tyr Pro Leu Met Ala Ile Ile Asn Pro Arg Arg Ser				
	500		505	